

# The cross-polytope, the ball, and the cube

Jordan Bell

jordan.bell@gmail.com

Department of Mathematics, University of Toronto

March 28, 2016

## 1 $\ell^q$ norms and volume of the unit ball

For  $x, y \in \mathbb{R}^n$ ,

$$\langle x, y \rangle = \sum_{j=1}^n x_j y_j.$$

Let  $e_1, \dots, e_n$  be the standard basis for  $\mathbb{R}^n$ .

$$x = \sum_{j=1}^n x_j e_j = \sum_{j=1}^n \langle x, e_j \rangle e_j.$$

For  $1 \leq q < \infty$  let

$$|x|_q = \left( \sum_{j=1}^n |x_j|^q \right)^{1/q}$$

and for  $q = \infty$  let

$$|x|_\infty = \max_{1 \leq j \leq n} |x_j|.$$

Then for  $1 \leq q \leq \infty$  let

$$B_q^n = \{x \in \mathbb{R}^n : |x|_q \leq 1\}.$$

For  $0 \leq k \leq n$  let  $\lambda_k$  be  $k$ -dimensional Lebesgue measure on  $\mathbb{R}^n$ . We calculate the volume of the unit ball with the  $\ell^q$  norm for  $1 \leq q < \infty$ .

**Theorem 1.** *For  $n \geq 1$  and for  $1 \leq q < \infty$ ,*

$$\lambda_n(B_q^n) = \frac{(2\Gamma(\frac{1}{q} + 1))^n}{\Gamma(\frac{n}{q} + 1)}.$$

*Proof.* For  $R \geq 0$  let  $V_q^n(R) = \lambda_n(R \cdot B_q^n)$ . For  $n = 1$ ,

$$V_q^1(R) = \lambda_1(R \cdot B_q^1) = \int_{-R \leq x_1 \leq R} d\lambda_1(x_1) = 2R.$$

By induction, suppose for some  $n$  that

$$V_q^n(R) = \frac{(2R\Gamma(\frac{1}{q} + 1))^n}{\Gamma(\frac{n}{q} + 1)}.$$

Using Fubini's theorem and the induction hypothesis and doing the change of variable  $x_{n+1} = Rt$  we calculate

$$\begin{aligned} V_q^{n+1}(R) &= \int_{|x_1|^q + \dots + |x_n|^q + |x_{n+1}|^q \leq R^q} d\lambda_{n+1}(x) \\ &= \int_{-R \leq x_{n+1} \leq R} \left( \int_{|x_1|^q + \dots + |x_n|^q \leq R^q - |x_{n+1}|^q} d\lambda_n(x_1, \dots, x_n) \right) d\lambda_1(x_{n+1}) \\ &= \int_{-R \leq x_{n+1} \leq R} V_q^n((R^q - |x_{n+1}|^q)^{1/q}) d\lambda_1(x_{n+1}) \\ &= \int_{-R \leq x_{n+1} \leq R} \frac{(2(R^q - |x_{n+1}|^q)^{1/q}\Gamma(\frac{1}{q} + 1))^n}{\Gamma(\frac{n}{q} + 1)} d\lambda_1(x_{n+1}) \\ &= \frac{(2\Gamma(\frac{1}{q} + 1))^n}{\Gamma(\frac{n}{q} + 1)} \int_{-1 \leq t \leq 1} (R^q - |Rt|^q)^{n/q} \cdot R d\lambda_1(t) \\ &= \frac{(2\Gamma(\frac{1}{q} + 1))^n}{\Gamma(\frac{n}{q} + 1)} \cdot R^{n+1} \cdot 2 \int_{0 \leq t \leq 1} (1 - t^q)^{n/q} d\lambda_1(t). \end{aligned}$$

Now, doing the change of variable  $u = t^q$ , namely  $t = u^{1/q}$  with  $t' = \frac{1}{q}u^{\frac{1}{q}-1}$  and using the beta function  $B(a, b) = \int_0^1 u^{a-1}(1-u)^{b-1} d\lambda_1(u)$ ,

$$\begin{aligned} \int_{0 \leq t \leq 1} (1 - t^q)^{n/q} d\lambda_1(t) &= \int_{0 \leq u \leq 1} (1 - u)^{n/q} \cdot \frac{1}{q}u^{\frac{1}{q}-1} d\lambda_1(u) \\ &= \frac{1}{q}B\left(\frac{1}{q}, \frac{n}{q} + 1\right). \end{aligned}$$

But  $B(a, b) = \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)}$ , and using  $\Gamma(a+1) = a\Gamma(a)$ ,

$$\frac{1}{q}B\left(\frac{1}{q}, \frac{n}{q} + 1\right) = \frac{1}{q} \frac{\Gamma\left(\frac{1}{q}\right)\Gamma\left(\frac{n}{q} + 1\right)}{\Gamma\left(\frac{1}{q} + \frac{n}{q} + 1\right)} = \frac{\Gamma\left(\frac{1}{q} + 1\right)\Gamma\left(\frac{n}{q} + 1\right)}{\Gamma\left(\frac{n+1}{q} + 1\right)}.$$

Therefore

$$\begin{aligned} V_q^{n+1}(R) &= \frac{(2\Gamma(\frac{1}{q} + 1))^n}{\Gamma(\frac{n}{q} + 1)} \cdot R^{n+1} \cdot 2 \cdot \frac{\Gamma\left(\frac{1}{q} + 1\right)\Gamma\left(\frac{n}{q} + 1\right)}{\Gamma\left(\frac{n+1}{q} + 1\right)} \\ &= \frac{(2R\Gamma(\frac{1}{q} + 1))^{n+1}}{\Gamma\left(\frac{n+1}{q} + 1\right)}, \end{aligned}$$

which proves the claim.  $\square$

$B_1^n$  is an  $n$ -dimensional **cross-polytope**,  $B_2^n$  is an  $n$ -dimensional **Euclidean ball**, and  $B_\infty^n$  is an  $n$ -dimensional **cube**.

$$\lambda_n(B_1^n) = \frac{2^n}{n!}, \quad \lambda_n(B_2^n) = \frac{\pi^{n/2}}{\Gamma(\frac{n}{2} + 1)}, \quad \lambda_n(B_\infty^n) = 2^n,$$

using  $\Gamma(n+1) = n!$  and  $\Gamma(\frac{3}{2}) = \frac{\sqrt{\pi}}{2}$ .

## 2 Intersection of a hyperplane and the cube

Let  $\xi \in S^{n-1}$  and  $t \in \mathbb{R}$ , and define

$$P_{\xi,t} = \{x \in \mathbb{R}^n : \langle x, \xi \rangle = t\}.$$

In particular,

$$\xi^\perp = P_{\xi,0}.$$

Let

$$A_\xi(t) = \lambda_{n-1}(P_{\xi,t} \cap B_\infty^n) = \int_{P_{\xi,t}} 1_{B_\infty^n}(x) d\lambda_{n-1}(x).$$

**Theorem 2.** For  $\xi \in S^{n-1}$  and  $t \in \mathbb{R}$ ,

$$A_\xi(t) = \frac{2^n}{\pi} \int_0^\infty \cos tr \cdot \prod_{i=1}^n \frac{2}{\xi_i r} \sin \xi_i r dr.$$

*Proof.* Then by Fubini's theorem,

$$\begin{aligned} \widehat{A}_\xi(\tau) &= \int_{\mathbb{R}} A_\xi(t) e^{-2\pi i t \tau} d\lambda_1(t) \\ &= \int_{\mathbb{R}} \left( \int_{P_{\xi,t}} 1_{B_\infty^n}(x) e^{-2\pi i \langle x, \xi \rangle \tau} d\lambda_{n-1}(x) \right) d\lambda_1(t) \\ &= \int_{\mathbb{R}^n} 1_{B_\infty^n}(x) e^{-2\pi i \langle x, \xi \rangle \tau} d\lambda_n(x). \end{aligned}$$

Now,

$$1_{B_\infty^n}(x) = (1_{B_\infty^1} \otimes \cdots \otimes 1_{B_\infty^1})(x) = \prod_{i=1}^n 1_{B_\infty^1}(x_i),$$

whence, by Fubini's theorem,

$$\begin{aligned} \int_{\mathbb{R}^n} 1_{B_\infty^n}(x) e^{-2\pi i \langle x, \xi \rangle \tau} d\lambda_n(x) &= \int_{\mathbb{R}^n} \left( \prod_{i=1}^n 1_{B_\infty^1}(x_i) e^{-2\pi i x_i \xi_i \tau} \right) d\lambda_n(x) \\ &= \prod_{i=1}^n \int_{\mathbb{R}} 1_{B_\infty^1}(x_i) e^{-2\pi i x_i \xi_i \tau} d\lambda_1(x_i). \end{aligned}$$

But, when  $\xi_i \tau \neq 0$ ,

$$\int_{\mathbb{R}} 1_{B_\infty^1}(x_i) e^{-2\pi i x_i \xi_i \tau} d\lambda_1(x_i) = \int_{-1}^1 e^{-2\pi i x_i \xi_i \tau} d\lambda_1(x_i) = \frac{1}{\pi \xi_i \tau} \sin 2\pi \xi_i \tau,$$

thus

$$\widehat{A}_\xi(\tau) = \prod_{i=1}^n \frac{1}{\pi \xi_i \tau} \sin 2\pi \xi_i \tau.$$

By the Fourier inversion theorem, using that  $\widehat{A}_\xi$  is an even function,

$$\begin{aligned} A_\xi(t) &= \int_{\mathbb{R}} \widehat{A}_\xi(\tau) e^{2\pi i t \tau} d\lambda_1(\tau) \\ &= \int_{\mathbb{R}} \widehat{A}_\xi(\tau) \cos 2\pi t \tau d\lambda_1(\tau) \\ &= 2 \int_0^\infty \widehat{A}_\xi(\tau) \cos 2\pi t \tau d\tau \\ &= 2 \int_0^\infty \cos 2\pi t \tau \cdot \prod_{i=1}^n \frac{1}{\pi \xi_i \tau} \sin 2\pi \xi_i \tau d\tau \\ &= \frac{2^n}{\pi} \int_0^\infty \cos t r \cdot \prod_{i=1}^n \frac{2}{\xi_i r} \sin \xi_i r dr. \end{aligned}$$

□

### 3 Schwartz functions

Let  $\mathcal{S}$  be the Fréchet space of Schwartz function  $\mathbb{R}^n \rightarrow \mathbb{C}$  and let  $\mathcal{S}'$  be the locally convex space of tempered distributions  $\mathcal{S} \rightarrow \mathbb{C}$ . If  $f : \mathbb{R}^n \rightarrow \mathbb{C}$  is locally integrable and there is some  $N$  such that

$$\int_{|x|_2 \leq R} |f(x)| d\lambda_n(x) = O(R^N), \quad R \rightarrow \infty,$$

it is a fact that

$$\phi \mapsto \langle f, \phi \rangle = \int_{\mathbb{R}^n} f(x) \phi(x) d\lambda_n(x), \quad \phi \in \mathcal{S},$$

is a tempered distribution.

**Lemma 3.** *For  $1 \leq q \leq \infty$  and for  $0 < h < n$ ,  $|x|_q^{-h}$  is a tempered distribution.*

*Proof.* For  $1 \leq q \leq 2$ ,

$$|x|_2 \leq |x|_q \leq n^{\frac{1}{q} - \frac{1}{2}} |x|_2,$$

and for  $2 \leq q \leq \infty$ ,

$$|x|_q \leq |x|_2 \leq n^{\frac{1}{2} - \frac{1}{q}} |x|_q.$$

Then for  $1 \leq q \leq 2$  and for  $0 < h < n$ , using polar coordinates and as  $\sigma(S^{n-1}) = \frac{2\pi^{(n+1)/2}}{\Gamma(\frac{n+1}{2})}$ ,

$$\begin{aligned} \int_{|x|_2 \leq R} |x|_q^{-h} d\lambda_n(x) &\leq \int_{|x|_2 \leq R} |x|_2^{-h} d\lambda_n(x) \\ &= \int_{S^{n-1}} \left( \int_0^\infty r^{-h} \cdot r^{n-1} dr \right) d\sigma \\ &= \frac{2\pi^{(n+1)/2}}{\Gamma(\frac{n+1}{2})} \cdot \int_0^R r^{-h+n-1} dr \\ &= \frac{2\pi^{(n+1)/2}}{\Gamma(\frac{n+1}{2})} \cdot \frac{r^{-h+n}}{-h+n} \Big|_0^R \\ &= \frac{2\pi^{(n+1)/2}}{\Gamma(\frac{n+1}{2})} \cdot \frac{R^{-h+n}}{-h+n} \\ &= O(R^{-h+n}). \end{aligned}$$

For  $2 \leq q \leq \infty$  and for  $0 < r < n$ ,

$$\begin{aligned} \int_{|x|_2 \leq R} |x|_q^{-h} d\lambda_n(x) &\leq \int_{|x|_2 \leq R} (n^{-\frac{1}{2} + \frac{1}{q}} |x|_2)^{-h} d\lambda_n(x) \\ &= n^{\frac{h}{2} - \frac{h}{q}} \int_{|x|_2 \leq R} |x|_2^{-h} d\lambda_n(x) \\ &= n^{\frac{h}{2} - \frac{h}{q}} \cdot \frac{2\pi^{(n+1)/2}}{\Gamma(\frac{n+1}{2})} \cdot \frac{R^{-h+n}}{-h+n} \\ &= O(R^{-h+n}). \end{aligned}$$

□

For  $\phi \in \mathcal{S}$  let

$$\widehat{\phi}(\xi) = \int_{\mathbb{R}^n} \phi(x) e^{-2\pi i \langle x, \xi \rangle} d\lambda_n(x).$$

For  $1 \leq q < \infty$  define  $c_q : \mathbb{R} \rightarrow \mathbb{R}$  by

$$c_q(z) = e^{-|z|^q}, \quad z \in \mathbb{R},$$

which belongs to  $\mathcal{S}(\mathbb{R})$ , and let  $\gamma_q = \widehat{c}_q$ .

For a tempered distribution  $T$ ,

$$\langle \widehat{T}, \phi \rangle = \langle T, \widehat{\phi} \rangle, \quad \phi \in \mathcal{S}.$$

Define  $f_{q,h}(x) = |x|_q^{-h}$ . We calculate the Fourier transform of the tempered distribution  $f_{q,h}$ .<sup>1</sup>

**Theorem 4.** Let  $0 < h < n$ . For  $1 \leq q < \infty$ ,

$$\widehat{f}_{q,h}(\xi) = \frac{q}{\Gamma(h/q)} \int_0^\infty t^{n-h-1} \prod_{j=1}^n \gamma_q(t\xi_j) dt,$$

and for  $q = \infty$ ,

$$\widehat{f}_{\infty,h}(\xi) = 2^n h \int_0^\infty t^{n-h-1} \prod_{j=1}^n \frac{\sin t\xi_j}{t\xi_j} dt$$

*Proof.* Suppose that  $1 \leq q < \infty$ . For  $x \neq 0$ , doing the change of variable  $z = t^{1/q}|x|_q^{-1}$ ,

$$\begin{aligned} \int_0^\infty z^{h-1} e^{-z^q|x|_q^q} dz &= \int_0^\infty (t^{1/q}|x|_q^{-1})^{h-1} e^{-t} |x|_q^{-1} \frac{1}{q} t^{\frac{1}{q}-1} dt \\ &= \frac{|x|_q^{-h}}{q} \int_0^\infty t^{\frac{h}{q}-1} e^{-t} dt \\ &= \frac{|x|_q^{-h}}{q} \cdot \Gamma(h/q), \end{aligned}$$

i.e.  $f_{q,h}(x) = \frac{q}{\Gamma(h/q)} \int_0^\infty z^{h-1} e^{-z^q|x|_q^q} dz$ .

For  $z > 0$  define  $F_{q,z} : \mathbb{R}^n \rightarrow \mathbb{R}$  by

$$F_{q,z}(x) = e^{-|zx|_q^q}, \quad x \in \mathbb{R}^n,$$

which is a Schwartz function. Doing the change of variable  $y = z \cdot x$  and using

---

<sup>1</sup>Alexander Koldobsky and Vladyslav Yaskin, *The Interface between Convex Geometry and Harmonic Analysis*, p. 9, Lemma 2.1.

Fubini's theorem,

$$\begin{aligned}
\widehat{F}_{q,z}(\xi) &= \int_{\mathbb{R}^n} e^{-|zx|_q^q} e^{-2\pi i \langle x, \xi \rangle} d\lambda_n(x) \\
&= \int_{\mathbb{R}^n} e^{-|y_1|^q - \dots - |y_n|^q} e^{-2\pi i \langle y, z^{-1}\xi \rangle} \cdot z^{-n} d\lambda_n(y) \\
&= z^{-n} \prod_{j=1}^n \int_{\mathbb{R}} e^{-|y_j|^q} e^{-2\pi i y_j \cdot z^{-1}\xi_j} d\lambda_1(y_j) \\
&= z^{-n} \prod_{j=1}^n \gamma_q(z^{-1}\xi_j).
\end{aligned}$$

Then for  $\phi \in \mathcal{S}$ ,

$$\begin{aligned}
\langle \widehat{f}_{q,h}, \phi \rangle &= \int_{\mathbb{R}^n} f_{q,h}(\xi) \widehat{\phi}(\xi) d\lambda_n(\xi) \\
&= \int_{\mathbb{R}^n} \left( \frac{q}{\Gamma(h/q)} \int_0^\infty z^{h-1} e^{-z^q |\xi|_q^q} dz \right) \widehat{\phi}(\xi) d\lambda_n(\xi) \\
&= \frac{q}{\Gamma(h/q)} \int_0^\infty z^{h-1} \left( \int_{\mathbb{R}^n} e^{-|z\xi|_q^q} \widehat{\phi}(\xi) d\lambda_n(\xi) \right) dz \\
&= \frac{q}{\Gamma(h/q)} \int_0^\infty z^{h-1} \langle F_{q,z}, \widehat{\phi} \rangle dz \\
&= \frac{q}{\Gamma(h/q)} \int_0^\infty z^{h-1} \langle \widehat{F}_{q,z}, \phi \rangle dz \\
&= \frac{q}{\Gamma(h/q)} \int_0^\infty z^{h-1-n} \left( \int_{\mathbb{R}^n} \prod_{j=1}^n \gamma_q(z^{-1}\xi_j) \cdot \phi(\xi) d\lambda_n(\xi) \right) dz \\
&= \int_{\mathbb{R}^n} \left( \frac{q}{\Gamma(h/q)} \int_0^\infty z^{h-1-n} \prod_{j=1}^n \gamma_q(z^{-1}\xi_j) dz \right) \phi(\xi) d\lambda_n(\xi).
\end{aligned}$$

This implies, doing the change of variable  $z = t^{-1}$ ,

$$\begin{aligned}
\widehat{f}_{q,h}(\xi) &= \frac{q}{\Gamma(h/q)} \int_0^\infty z^{h-1-n} \prod_{j=1}^n \gamma_q(z^{-1}\xi_j) dz \\
&= \frac{q}{\Gamma(h/q)} \int_0^\infty t^{n-h-1} \prod_{j=1}^n \gamma_q(t\xi_j) dt.
\end{aligned}$$

□

## 4 $\gamma_q$

We remind ourselves that  $c_q(z) = e^{-|z|^q}$ ,  $z \in \mathbb{R}$ , and  $\gamma_q = \widehat{c}_q$ . We prove that  $\gamma_q$  is positive and logconvex.<sup>2</sup>

**Theorem 5.** *For  $1 \leq q \leq 2$ ,  $\gamma_q(\sqrt{z}) > 0$ , and  $z \mapsto \log \gamma_q(\sqrt{z})$  is convex on  $\mathbb{R}_{\geq 0}$ .*

*Proof.* Let  $0 < \alpha \leq 1$ , and for  $z \in [0, \infty)$  let  $f(z) = \exp z$  and  $g(z) = -z^\alpha$ . Then for  $k \in \mathbb{Z}_{\geq 0}$  and  $z \in (0, \infty)$ ,

$$g^{(k)}(z) = -k! \binom{\alpha}{k} z^{\alpha-k}, \quad \operatorname{sgn} g^{(k)}(z) = (-1)^k.$$

For  $n \geq 1$ , **Faà di Bruno's formula** tells us

$$(f \circ g)^{(n)}(z) = \sum_{(m_1, \dots, m_n), 1 \cdot m_1 + \dots + n \cdot m_n = n} \frac{n!}{m_1! \cdots m_n!} (f^{(m_1 + \dots + m_n)} \circ g)(z) \\ \cdot \prod_{k=1}^n \left( \frac{g^{(k)}(z)}{k!} \right)^{m_k}.$$

Then

$$(-1)^n (f \circ g)^{(n)}(z) = \sum_{(m_1, \dots, m_n), 1 \cdot m_1 + \dots + n \cdot m_n = n} \frac{n!}{m_1! \cdots m_n!} (\exp \circ g)(z) \\ \cdot \prod_{k=1}^n \left( (-1)^k \frac{g^{(k)}(z)}{k!} \right)^{m_k} \\ \geq 0.$$

This shows that  $f \circ g$  is **completely monotone**. Furthermore,  $(f \circ g)(0) = 1$ , so by the **Bernstein-Widder theorem** there is a Borel probability measure  $\mu$  on  $[0, \infty)$  such that<sup>3</sup>

$$(f \circ g)(z) = \int_{[0, \infty)} e^{-zt} d\mu(t), \quad z \in [0, \infty).$$

With  $\alpha = \frac{q}{2}$ , there is thus a Borel probability measure  $\mu_q$  on  $[0, \infty)$  such that

$$\exp(-z^{q/2}) = \int_{[0, \infty)} e^{-zt} d\mu_q(t), \quad z \in [0, \infty).$$

Then for  $z \in \mathbb{R}$ ,

$$c_q(z) = \exp(-|z|^q) = \exp(-(z^2)^{q/2}) = \int_{[0, \infty)} e^{-z^2 t} d\mu_{q/2}(t).$$

---

<sup>2</sup>Alexander Koldobsky and Vladyslav Yaskin, *The Interface between Convex Geometry and Harmonic Analysis*, p. 4, Lemma 1.4.

<sup>3</sup><http://individual.utoronto.ca/jordanbell/notes/completelymonotone.pdf>

For  $w \in \mathbb{R}$  we calculate, using the Fourier transform of a Gaussian,<sup>4</sup>

$$\begin{aligned}
\gamma_q(w) &= \int_{\mathbb{R}} e^{-2\pi i w z} c_q(z) d\lambda_1(z) \\
&= \int_{\mathbb{R}} e^{-2\pi i w z} \left( \int_{[0, \infty)} e^{-z^2 t} d\mu_{q/2}(t) \right) d\lambda_1(z) \\
&= \int_{[0, \infty)} \left( \int_{\mathbb{R}} e^{-tz^2} e^{-2\pi i w z} d\lambda_1(z) \right) d\mu_{q/2}(t) \\
&= \int_{[0, \infty)} \sqrt{\frac{\pi}{t}} \exp\left(-\frac{(\pi w)^2}{t}\right) d\mu_{q/2}(t) \\
&= \pi^{1/2} \int_{[0, \infty)} t^{-1/2} e^{-\pi^2 w^2 / t} d\mu_{q/2}(t).
\end{aligned}$$

From the final expression it is evident that  $\gamma_k(w) > 0$ . Furthermore, for  $w_1, w_2 \in (0, \infty)$ , using the Cauchy-Schwarz inequality,

$$\begin{aligned}
\log \gamma_q \left( \sqrt{\frac{w_1 + w_2}{2}} \right) &= \frac{1}{2} \log \left( \pi^{1/2} \int_{[0, \infty)} t^{-1/4} e^{-\pi^2 \cdot \frac{w_1}{2t}} \cdot t^{-1/4} e^{-\pi^2 \cdot \frac{w_2}{2t}} d\mu_{q/2}(t) \right)^2 \\
&\leq \frac{1}{2} \log \left( \pi \int_{[0, \infty)} t^{-1/2} e^{-\pi^2 \cdot \frac{w_1}{t}} d\mu_{q/2}(t) \right. \\
&\quad \cdot \left. \int_{[0, \infty)} t^{-1/2} e^{-\pi^2 \cdot \frac{w_2}{t}} d\mu_{q/2}(t) \right) \\
&= \frac{1}{2} \log \left( \pi^{1/2} \int_{[0, \infty)} t^{-1/2} e^{-\pi^2 \cdot \frac{w_1}{t}} d\mu_{q/2}(t) \right) \\
&\quad + \frac{1}{2} \log \left( \pi^{1/2} \int_{[0, \infty)} t^{-1/2} e^{-\pi^2 \cdot \frac{w_2}{t}} d\mu_{q/2}(t) \right) \\
&= \frac{1}{2} \log \gamma_q(\sqrt{w_1}) + \frac{1}{2} \log \gamma_q(\sqrt{w_2}).
\end{aligned}$$

Because  $w \mapsto \log \gamma_q(\sqrt{w})$  is continuous, this suffices to prove that it is convex.  $\square$

---

<sup>4</sup><http://individual.utoronto.ca/jordanbell/notes/gaussianintegrals.pdf>