

# Gibbs measures and the Ising model

Jordan Bell

July 1, 2014

Let  $\Lambda$  be a finite subset of  $\mathbb{Z}^2$  and let  $\Lambda' = \mathbb{Z}^2 \setminus \Lambda$ . Let  $\sigma' \in \{-1, +1\}^{\Lambda'}$ , a fixed configuration of spins outside  $\Lambda$ . Let  $\Omega = \{-1, +1\}^\Lambda$ ;  $\Omega$  is the space of all configurations of spins on  $\Lambda$ . We define a Hamiltonian  $H_\Lambda(\cdot|\sigma') : \Omega \rightarrow \mathbb{R}$  (depending on the fixed external configuration  $\sigma'$ ) by

$$H_\Lambda(\sigma|\sigma') = - \sum_{\substack{x, y \in \Lambda \\ |x-y|=1}} \sigma(x)\sigma(y) - \sum_{\substack{x \in \Lambda, y \in \Lambda' \\ |x-y|=1}} \sigma(x)\sigma'(y).$$

$H_\Lambda(\cdot|\sigma')$  gives the energy of a configuration  $\sigma \in \Omega$ , conditioned on the external configuration  $\sigma'$ .

For a parameter  $\beta > 0$  (called the *inverse temperature*), we define the *partition function* by

$$Z(\beta, \Lambda, \sigma') = \sum_{\sigma \in \Omega} \exp(-\beta H_\Lambda(\sigma|\sigma')).$$

Then we define the *Gibbs distribution* for the configuration space  $\Omega$ , depending on the external configuration  $\sigma'$ , by

$$P_{\beta, \Lambda}(\sigma|\sigma') = \frac{1}{Z(\beta, \Lambda, \sigma')} \exp(-\beta H(\sigma|\sigma')).$$

The purpose of the partition function is to normalize the above expression to be a probability measure on the configuration space  $\Omega$ .

For example, let  $\Lambda$  be a square of side length 3 centred at the origin, and take  $\sigma'$  to be an external configuration of all negative spins. Define  $\sigma \in \Omega$  by

$$\begin{array}{lll} \sigma(-1, 1) = +1 & \sigma(0, 1) = +1 & \sigma(1, 1) = -1 \\ \sigma(-1, 0) = -1 & \sigma(0, 0) = +1 & \sigma(1, 0) = -1 \\ \sigma(-1, -1) = -1 & \sigma(0, -1) = -1 & \sigma(1, -1) = +1. \end{array}$$

We show this configuration in Figure 1. We calculate that the energy of this configuration is  $H_\Lambda(\sigma|\sigma') = 0$ . We can calculate the energy of this configuration in a different way, using line segments separating lattice points with different spins, as follows. For an  $n \times n$  square, there are  $2n(n+1)$  nearest neighbor interactions. Put a line segment between every two lattice points with different spins; let  $B(\sigma|\sigma')$  be the set of these line segments. We show this in Figure 2.




Figure 1: An example of a configuration (and negative external spins)




Figure 2: Calculating energy using contours

Generally, if  $\Lambda$  is an  $n \times n$  square then we have

$$H_\Lambda(\sigma|\sigma') = -2n(n+1) + 2|B(\sigma|\sigma')|.$$

Indeed, in our above example,  $n = 3$  and  $|B(\sigma|\sigma')| = 12$ , so the above expression is  $-24 + 2 \cdot 12 = 0$ , and we have already calculated that  $H_\Lambda(\sigma|\sigma') = 0$ . What matters is that if we know the external configuration, then to describe the configuration inside a region  $\Lambda$  it suffices to know the edges that separate opposite spins. And since the energy of any configuration has the term  $-2n(n+1)$  and this appears in the numerator and denominator of the expression for the Gibbs distribution, we can omit it to calculate the Gibbs distribution. By a *contour* we mean a closed path of edges that does not intersect itself. We can express the Gibbs distribution in terms of contours as

$$P_{\beta,\Lambda}(\sigma|\sigma') = \frac{\prod_{\gamma \in \Gamma(\sigma,\sigma')} \exp(-2|\gamma|)}{\sum_{\Gamma} \prod_{\gamma \in \Gamma} \exp(-2\beta|\gamma|)};$$

$\Gamma(\sigma,\sigma')$  is the set of contours corresponding to the configuration  $\sigma$  with the external configuration  $\sigma'$ , and the summation is over all sets  $\Gamma$  of nonintersecting contours.

We are not in fact interested in the Gibbs distribution on the configurations on a finite subset  $\Lambda$  of  $\mathbb{Z}^2$ , but instead limits of Gibbs distributions with  $\Lambda_n \rightarrow \mathbb{Z}^2$ . A Gibbs distribution  $P_{\beta,\Lambda}(\cdot|\sigma')$  on  $\Omega$  is in fact a probability measure on  $\{+1, -1\}^{\mathbb{Z}^2}$ : for  $\sigma \in \{+1, -1\}^{\mathbb{Z}^2}$ , a configuration on the plane, we define

$$\tilde{P}_{\beta,\Lambda}(\sigma|\sigma') = \begin{cases} 0 & \sigma|_{\Lambda'} \neq \sigma' \\ P_{\beta,\Lambda}((\sigma|_{\Lambda})|\sigma') & \sigma|_{\Lambda'} = \sigma'. \end{cases}$$

Fix some  $\beta$ . Let  $\Lambda_n$  be a sequence of  $n \times n$  squares centred at the origin, let  $\sigma'_{n,+}$  be a sequence of external configurations where all lattice points outside  $\Lambda_n$  have positive spins, and let  $\sigma'_{n,-}$  be a sequence of external configurations where all lattice points outside  $\Lambda_n$  have negative spins. Let  $P_{n,+}$  be the sequence of Gibbs distributions corresponding to the positive external spins, and let  $P_{n,-}$  be the sequence of Gibbs distributions corresponding to the negative external spins. These extend to probability measures  $\tilde{P}_{n,+}$  and  $\tilde{P}_{n,-}$  on  $\{+1, -1\}^{\mathbb{Z}^2}$ . Since  $\{+1, -1\}$  is a compact metrizable space, the product  $\{+1, -1\}^{\mathbb{Z}^2}$  is a compact metrizable space and thus the space of probability measures on it is compact. Hence the sequence  $\tilde{P}_{n,+}$  has at least one limit point, say  $P_+$ , and the sequence  $\tilde{P}_{n,-}$  has at least one limit point, say  $P_-$ . We shall show that  $P_+ \neq P_-$ , namely that there is not a unique limit Gibbs measure on the set of all configurations on  $\mathbb{Z}^2$ .

Let  $V_+ = \{\sigma \in \{+1, -1\}^{\mathbb{Z}^2} : \sigma(0) = +1\}$  and  $V_- = \{\sigma \in \{+1, -1\}^{\mathbb{Z}^2} : \sigma(0) = -1\}$ . Suppose that for all  $n$  we had  $\tilde{P}_{n,+}(V_-) < \frac{1}{3}$ . Taking limits we have that  $P_+(V_-) \leq \frac{1}{3}$  and so  $P_+(V_+) \geq \frac{2}{3}$  (since the events  $V_+$  and  $V_-$  are disjoint and their union is the set of all configurations on  $\mathbb{Z}^2$ ). But  $\tilde{P}_{n,+}(V_-) = \tilde{P}_{n,-}(V_+)$ , so taking limits we also get  $P_-(V_+) \leq \frac{1}{3}$ . Therefore the measures  $P_+$  and  $P_-$

give different measures to the set  $V_+$ , so they are distinct. Thus to show that the measures  $P_+$  and  $P_-$  are distinct it suffices to show that for all  $n$  we have  $\tilde{P}_{n,+}(V_-) < \frac{1}{3}$ .

We have

$$\begin{aligned} \tilde{P}_{n,+}(V_-) &\leq \text{Prob}(\text{there exists a contour } \gamma \subset B(\sigma|\sigma'), 0 \in \text{Int}(\gamma)) \\ &\leq \sum_{0 \in \text{Int}(\gamma)} \text{Prob}(\gamma \subset B(\sigma|\sigma')) \\ &\leq \sum_{0 \in \text{Int}(\gamma)} \exp(-2\beta|\gamma|). \end{aligned}$$

The above sum is over all contours such that the origin lies in their interior. We can write the set of all contours around the origin as a union of the set of all contours of length  $k$  around the origin,  $k \geq 4$ . There are at most  $(\frac{k}{4})^2 4^k$  contours of length  $k$  around the origin. Therefore

$$\tilde{P}_{n,+}(V_-) \leq \sum_{k=4}^{\infty} \frac{k^2}{16} \cdot 4^k \exp(-2\beta k).$$

As  $\beta \rightarrow \infty$ , this is  $O(\exp(-8\beta))$ . In particular there is some  $\beta_0$  such that if  $\beta \geq \beta_0$  then for all  $n$  we have  $\tilde{P}_{n,+}(V_-) < \frac{1}{3}$ . This shows that the limit Gibbs measures gives different measures to the set  $V_+$ , hence they are distinct.

## Further reading

Minlos [4], Sinai [6], Cipra [1], Simon [5], Le Ny [3], Kadanoff [2].

## References

- [1] Barry A. Cipra, *An introduction to the Ising model*, Amer. Math. Monthly **94** (1987), no. 10, 937–959.
- [2] Leo P. Kadanoff, *More is the same; phase transitions and mean field theories*, J. Stat. Phys. **137** (2009), 777–797.
- [3] Arnaud Le Ny, *Introduction to (generalized) Gibbs measures*, 2007, arXiv:0712.1171.
- [4] R. A Minlos, *Introduction to mathematical statistical physics*, University Lecture Series, vol. 19, American Mathematical Society, Providence, R.I., 2000.
- [5] Barry Simon, *The statistical mechanics of lattice gases*, vol. I, Princeton University Press, 1993.

- [6] Ya. G. Sinai, *Theory of phase transitions: rigorous results*, Pergamon Press, Oxford, 1982, Translated from the Russian.