

Trace class operators and Hilbert-Schmidt operators

Jordan Bell

jordan.bell@gmail.com

Department of Mathematics, University of Toronto

April 18, 2016

1 Introduction

If X, Y are normed spaces, let $\mathcal{B}(X, Y)$ be the set of all bounded linear maps $X \rightarrow Y$. If $T : X \rightarrow Y$ is a linear map, I take it as known that T is bounded if and only if it is continuous if and only if $E \subseteq X$ being bounded implies that $T(E) \subseteq Y$ is bounded. I also take it as known that $\mathcal{B}(X, Y)$ is a normed space with the operator norm, that if Y is a Banach space then $\mathcal{B}(X, Y)$ is a Banach space, that if X is a Banach space then $\mathcal{B}(X) = \mathcal{B}(X, X)$ is a Banach algebra, and that if H is a Hilbert space then $\mathcal{B}(H)$ is a C^* -algebra. An *ideal* I of a Banach algebra is an ideal of the algebra: to say that I is an ideal does not demand that I is a Banach subalgebra, i.e. does not demand that I is a closed subset of the Banach algebra. I is a **-ideal* of a C^* -algebra if I is an ideal of the algebra and if $A \in I$ implies that $A^* \in I$.

If X and Y are normed spaces, we take as known that the *strong operator topology* on $\mathcal{B}(X, Y)$ is coarser than the norm topology on $\mathcal{B}(X, Y)$, and thus if $T_n \rightarrow T$ in the operator norm, then $T_n \rightarrow T$ in the strong operator topology.

If X is a normed space, M is a dense subspace of X , Y is a Banach space and $T : M \rightarrow Y$ is a bounded linear operator, then there is a unique element of $\mathcal{B}(X, Y)$ whose restriction to M is equal to T , and we also denote this by T .¹

If X is a normed space, define $\langle \cdot, \cdot \rangle : X \times X^* \rightarrow \mathbb{C}$ by

$$\langle x, \lambda \rangle = \lambda(x), \quad x \in X, \lambda \in X^*.$$

This is called the *dual pairing*. If X and Y are normed spaces and $T \in \mathcal{B}(X, Y)$, it can be proved that there is a unique $T^* \in \mathcal{B}(Y^*, X^*)$ such that

$$\langle Tx, \lambda \rangle = \langle x, T^* \lambda \rangle, \quad x \in X, \lambda \in Y^*,$$

called the *adjoint* of T , and that the adjoint satisfies $\|T\| = \|T^*\|$.²

¹This is an instance of a result about topological vector spaces and Fréchet spaces. See Walter Rudin, *Functional Analysis*, second ed., p. 40, chapter 1, ex. 19.

²Walter Rudin, *Functional Analysis*, second ed., p. 98, Theorem 4.10.

I give precise statements of any statement that I want to use without proof. If a fact is not straightforward to prove and is not easy to look up (perhaps because it does not have a standardized name), I give a citation to a statement that is the precise version I use. I am making a point to write out full proofs of some tedious but essential arguments about Hilbert-Schmidt operators and trace class operators.³

2 Finite rank operators

In this section, X and Y denote Banach spaces. We say that a linear map $T : X \rightarrow Y$ has *finite rank* if $T(X)$ is a finite dimensional subspace of Y . A finite rank operator need not be bounded: If X is infinite dimensional, let \mathcal{E} be a Hamel basis for X and let $\{e_n : n \in \mathbb{N}\}$ be a subset of this basis. Defining a linear map on a Hamel basis determines it on the vector space. Define $T : X \rightarrow \mathbb{C}$ by $Te_n = n\|e_n\|$ and $Te = 0$ if e is not in the countable subset. Then T is not bounded; but its range has dimension 1, so T has finite rank. We define $\mathcal{B}_{00}(X, Y)$ to be the set of $T \in \mathcal{B}(X, Y)$ that have finite rank. It is apparent that $\mathcal{B}_{00}(X, Y)$ is a vector space.

If $f \in X^*$ and $y \in Y$, we define $y \otimes f : X \rightarrow Y$ by

$$y \otimes f(x) = f(x)y.$$

$y \otimes f : X \rightarrow Y$ is linear, and

$$\|y \otimes f\| = \sup_{\|x\| \leq 1} \|(y \otimes f)(x)\| = \sup_{\|x\| \leq 1} \|f(x)y\| = \|y\| \sup_{\|x\| \leq 1} |f(x)| \leq \|y\| \sup_{\|x\| \leq 1} \|f\| \|x\|,$$

so $\|y \otimes f\| \leq \|f\| \|y\|$, so $y \otimes f$ is bounded. $(y \otimes f)(H) \subseteq \text{span}\{y\}$, so $y \otimes f$ has finite rank. Therefore $y \otimes f \in \mathcal{B}_{00}(X, Y)$.

The following theorem gives a representation for bounded finite rank operators.⁴

Theorem 1. *If $T \in \mathcal{B}_{00}(X, Y)$ and w_1, \dots, w_k is a basis for $T(X)$, then there are unique $f_1, \dots, f_k \in X^*$ such that*

$$T = \sum_{j=1}^k w_j \otimes f_j.$$

³Many of the proofs that I give are vastly expanded from what is written in the references I used; I simply decided to write down every step that I did, and thus a reader should be able to read this note without needing having to work out calculations on paper or without realizing partway through that I was tacitly identifying things or that what I said is true only under conditions I left unstated because I thought them too obvious. Indeed there is no royal road through mathematics, but we do not need to break up the asphalt and destroy the signage to make travelling what roads there are a trial of one's skill.

⁴Y. A. Abramovich and C. D. Aliprantis, *An Invitation to Operator Theory*, p. 124, Lemma 4.2. In this reference, what we write as $y \otimes f$ they write as $f \otimes y$.

For $y \in Y$, define $F_y : Y^* \rightarrow \mathbb{C}$ by $F_y(\lambda) = \lambda(y)$. One checks that $F_y \in (Y^*)^*$. Often one writes $y = F_y$, which is fine as long as we keep in mind whether we are using y as an element of Y or as an element of $(Y^*)^*$.

Theorem 2. *If there are $w_1, \dots, w_k \in Y$ and $f_1, \dots, f_k \in X^*$ such that*

$$T = \sum_{j=1}^k w_j \otimes f_j,$$

then

$$T^* = \sum_{j=1}^k f_j \otimes w_j.$$

Proof. Let $y \in Y$ and $f \in X^*$. If $x \in X$ and $g \in Y^*$, then

$$\langle (y \otimes f)x, g \rangle = \langle f(x)y, g \rangle = f(x)\langle y, g \rangle = f(x)g(y),$$

where $\langle \cdot, \cdot \rangle : Y \times Y^* \rightarrow \mathbb{C}$ is the dual pairing. But

$$f(x)g(y) = \langle x, g(y)f \rangle = \langle x, F_y(g)f \rangle = \langle x, (F_y \otimes f)(g) \rangle.$$

where $\langle \cdot, \cdot \rangle : X \times X^* \rightarrow \mathbb{C}$ is the dual pairing. We have $F_y \otimes f \in \mathcal{B}(Y^*, X^*)$, and hence

$$(y \otimes f)^* = F_y \otimes f = y \otimes f.$$

This shows that the adjoint of each term $w_j \otimes f_j$ in T is $f_j \otimes w_j$, and the adjoint of a sum is the sum of the adjoints of the terms, completing the proof. \square

The above two theorems together show that if $T \in \mathcal{B}_{00}(X, Y)$ then $T^* \in \mathcal{B}_{00}(Y^*, X^*)$.

Theorem 3. *If X is a Banach space, then $\mathcal{B}_{00}(X)$ is a two sided ideal in the Banach algebra $\mathcal{B}(X)$.*

Proof. We have stated already that $\mathcal{B}_{00}(X, Y)$ is a vector space, and here $Y = X$. If $A \in \mathcal{B}_{00}(X)$ and $T \in \mathcal{B}(X)$, then $AT \in \mathcal{B}(X)$, and

$$A(T(X)) \subseteq A(X),$$

which is finite dimensional, so $AT \in \mathcal{B}_{00}(X)$. $TA \in \mathcal{B}(X)$, and $T(A(X))$ is the image of a finite dimensional subspace under T , and so is itself finite dimensional. Hence $TA \in \mathcal{B}_{00}(X)$. \square

3 Compact operators

We say that a metric space M is *totally bounded* if for every ϵ there are finitely many balls of radius ϵ whose union equals M . The Heine-Borel theorem states that a metric space is compact if and only if it is complete and totally bounded. If S is a subset of a complete metric space M and \overline{S} is compact, then by the Heine-Borel theorem it is totally bounded, and any subset of a totally bounded metric space is itself a totally bounded metric space, so S is totally bounded. On the other hand, if $S \subseteq M$ is totally bounded, then one proves that \overline{S} is also totally bounded. As \overline{S} is a closed subset of the complete metric space M , \overline{S} is a complete metric space, and hence by the Heine-Borel theorem it is compact. We say that a subset of a metric space is *precompact* if its closure is compact, and thus a subset of a complete metric space is precompact if and only if it is totally bounded.

Let X and Y be Banach spaces. If $T : X \rightarrow Y$ is a linear map and U is the open unit ball in X , we say that T is *compact* if the closure of $T(U)$ in Y is compact. Therefore, to say that a linear map $T : X \rightarrow Y$ is compact is to say that $T(U)$ is totally bounded.

It doesn't take long to show that if $T : X \rightarrow Y$ is linear and compact then it is bounded, so there is no difference between stating that something is a bounded compact operator and stating that it is a compact operator. The following is often a convenient characterization of a compact operator.

Theorem 4. *A linear map $T : X \rightarrow Y$ is compact if and only if for every bounded sequence $x_n \in X$ there is a subsequence $x_{a(n)}$ such that $Tx_{a(n)}$ converges in Y .*

We denote the set of compact operators $X \rightarrow Y$ by $\mathcal{B}_0(X, Y)$. It is apparent that $\mathcal{B}_0(X, Y)$ is a vector space. For $T \in \mathcal{B}(X, Y)$, it is a fact that $T \in \mathcal{B}_0(X, Y)$ if and only if $T^* \in \mathcal{B}_0(Y^*, X^*)$.⁵

The following theorem states if a sequence of compact operators converges to a bounded operator, then that operator is compact.⁶ Since $\mathcal{B}(X, Y)$ is a Banach space, this implies that $\mathcal{B}_0(X, Y)$ is a Banach space with the operator norm.

Theorem 5. *If X and Y are Banach spaces, then $\mathcal{B}_0(X, Y)$ is a closed subspace of the Banach space $\mathcal{B}(X, Y)$.*

The following theorem shows that a bounded finite rank operator is a compact operator. Since a limit of compact operators is a compact operator, it follows from this that a limit of bounded finite rank operators is a compact operator.

Theorem 6. *If $T \in \mathcal{B}_{00}(X, Y)$ then $T \in \mathcal{B}_0(X, Y)$.*

⁵Walter Rudin, *Functional Analysis*, second ed., p. 105, Theorem 4.19.

⁶Walter Rudin, *Functional Analysis*, second ed., p. 104, Theorem 4.18 (c).

Proof. Let U be the open unit ball in X . Since T is bounded and U is a bounded set in X , $T(U)$ is a bounded set in Y . But $T(X)$ is finite dimensional, so $T(U)$ is a bounded set in a finite dimensional vector space and hence by the Heine-Borel theorem, the closure of $T(U)$ in $T(X)$ is a compact subset of $T(X)$. $T(X)$ is finite dimensional so is a closed subset of Y , and hence the closure of $T(U)$ in $T(X)$ is equal to the closure of $T(U)$ in Y . Hence the closure of $T(U)$ in Y is a compact subset of Y . (If E is a topological space and $C \subseteq D \subseteq E$, then the subspace topology C inherits from E is the same as the subspace topology it inherits from D , so if $K \subseteq T(X) \subseteq Y$ then to say that K is compact in $T(X)$ is equivalent to saying that K is compact in Y .) \square

Theorem 7. *If X is a Banach space, then $\mathcal{B}_0(X)$ is a two sided ideal in the Banach algebra $\mathcal{B}(X)$.*

Proof. We have stated already that $\mathcal{B}_0(X, Y)$ is a closed subspace of $\mathcal{B}(X, Y)$, and here $Y = X$. Let $K \in \mathcal{B}_0(X)$ and $T \in \mathcal{B}(X)$. On the one hand, if x_n is a bounded sequence in X , then $\|Tx_n\| \leq \|T\|\|x_n\|$, so Tx_n is a bounded sequence in X . Hence there is a subsequence $Tx_{a(n)}$ such that $K(Tx_{a(n)}) = (KT)x_{a(n)}$ converges, showing that KT is compact.

On the other hand, if x_n is a bounded sequence in X , then there is a subsequence $x_{a(n)}$ such that $Kx_{a(n)}$ converges to some x . T is continuous, so $T(Kx_{a(n)})$ converges to Tx , showing that TK is compact. \square

If X is not separable, then the image of the identity map id_X is not separable, so the image of a bounded linear operator need not be separable. However, the following theorem shows that the image of a compact operator is separable. Check that if a subset of a metric space is separable then its closure is separable. From this and Theorem 8 we get that the closure of the image of a compact operator is separable.

Theorem 8. *If $K \in \mathcal{B}_0(X, Y)$, then $K(X)$ with the subspace topology from Y is separable.*

Proof. Let $U_n = \{x \in X : \|x\| < n\}$. Then $\overline{K(U_n)}$ is compact. It is a fact that a compact metric space is separable, hence $\overline{K(U_n)}$ is separable. A subset of a separable metric space is itself separable with the subspace topology, so let L_n be a countable dense subset of $K(U_n)$. Let $L = \bigcup_{n=1}^{\infty} L_n$, which is countable. It is straightforward to verify that L is a dense subset of

$$K(X) = \bigcup_{n=1}^{\infty} K(U_n).$$

Therefore, $K(X)$ is separable. \square

4 Hilbert spaces

We showed that if X is a Banach space then both $\mathcal{B}_{00}(X)$ and $\mathcal{B}_0(X)$ are two sided ideals in the Banach algebra $\mathcal{B}(X)$. We also showed that if $A \in \mathcal{B}_{00}(X)$

then $A^* \in \mathcal{B}_{00}(X^*)$, and that if $A \in \mathcal{B}_0(X)$ then $A^* \in \mathcal{B}_0(X^*)$. If H is a Hilbert space then $\mathcal{B}(H)$ is a C^* -algebra (as the adjoint of $A \in \mathcal{B}(H)$ is not just an element of $\mathcal{B}(H^*)$, but can be identified with an element of $\mathcal{B}(H)$) and what we have shown implies that $\mathcal{B}_{00}(H)$ and $\mathcal{B}_0(H)$ are two sided $*$ -ideals in the C^* -algebra $\mathcal{B}(H)$.

If $S_\alpha, \alpha \in I$ are subsets of a Hilbert space H , we denote by $\bigvee_{\alpha \in I} S_\alpha$ the closure of the span of $\bigcup_{\alpha \in I} S_\alpha$. If S_1, S_2 are subsets of H , we write $S_1 \perp S_2$ if for every $s_1 \in S_1$ and $s_2 \in S_2$ we have $\langle s_1, s_2 \rangle = 0$. If V is a closed subspace of H , then $H = V \oplus V^\perp$, and the *orthogonal projection* onto V is the map $P : H \rightarrow H$ defined by $P(v + w) = v$ for $v \in V, w \in V^\perp$. $P(H) = V$, so rather than first fixing a closed subspace and then talking about the orthogonal projection onto that subspace, one often speaks about an orthogonal projection, which is the orthogonal projection onto its image. It is straightforward to check that if an orthogonal projection is nonzero then it has norm 1, and that an orthogonal projection is a positive operator.

The following theorem is an explicit version of Theorem 1 for orthogonal projections.

Theorem 9. *Let $\{e_1, \dots, e_n\}$ be orthonormal in H and let $M = \bigvee_{k=1}^n \{e_k\}$. If P is the orthogonal projection onto M , then*

$$P = \sum_{k=1}^n e_k \otimes e_k, \quad Ph = \sum_{k=1}^n \langle h, e_k \rangle e_k, \quad h \in H.$$

Proof. As M is finite dimensional it is closed, and hence $H = M \oplus M^\perp$. Let $h = h_1 + h_2$, $h_1 \in M, h_2 \in M^\perp$; as P is the orthogonal projection onto M , we have $Ph = h_1$.

Let $Qh = \sum_{k=1}^n \langle h, e_k \rangle e_k$. We have to show that $Qh = Ph$. For each $1 \leq j \leq n$, using that $\langle e_k, e_j \rangle = \delta_{k,j}$ we get

$$\langle Qh, e_j \rangle = \sum_{k=1}^n \langle h, e_k \rangle \langle e_k, e_j \rangle = \langle h, e_j \rangle.$$

Hence, if $1 \leq j \leq n$ then $\langle h - Qh, e_j \rangle = 0$. As $\{e_j\}$ are an orthonormal basis for M , this implies that $h - Qh \perp M$, and so $h - Qh \in M^\perp$. That is,

$$h_1 + h_2 - Qh \in M^\perp,$$

and as $h_2 \in M^\perp$ it follows that $h_1 - Qh \in M^\perp$. But $h_1 - Qh \in M$ ($h_1 \in M$ by definition, and Qh is a sum of elements in M), so $h_1 - Qh = 0$. As $Ph = h_1$, this means that $Ph = Qh$, completing the proof. \square

We say that a Banach space X has the *approximation property* if for each $A \in \mathcal{B}_0(X)$ there is a sequence $A_n \in \mathcal{B}_{00}(X)$ such that $A_n \rightarrow A$. A result of Per Enflo shows that there are Banach spaces that do not have the approximation property. However, in the following theorem we show that every Hilbert space has the approximation property.

Theorem 10. *If H is a Hilbert space and $A \in \mathcal{B}_0(H)$, then there is a sequence $A_n \in \mathcal{B}_0(H)$ such that $A_n \rightarrow A$.*

Proof. By Theorem 8, $V = \overline{A(H)}$ is separable. V is a closed subspace of the Hilbert space H , so is itself a Hilbert space. If V has finite dimension then A is itself finite rank. Otherwise, let $\{e_n : n \geq 1\}$ be an orthonormal basis for V , and let P_n be the orthogonal projection onto $\bigvee_{j=1}^n \{e_j\}$. $P_n \in \mathcal{B}_0(H)$, and define $A_n = P_n A \in \mathcal{B}_0(H)$.

In any Hilbert space, if v_α is an orthonormal basis then $\text{id}_H = \sum_\alpha v_\alpha \otimes v_\alpha$, where the series converges in the strong operator topology (see §5). Thus, for any $h \in V$ we have

$$Ah = \sum_{k=1}^{\infty} \langle Ah, e_k \rangle e_k,$$

where the series converges in H . By Theorem 9,

$$A_n h = P_n Ah = \sum_{k=1}^n \langle Ah, e_k \rangle e_k,$$

and therefore $\|A_n h - Ah\| \rightarrow 0$ as $n \rightarrow \infty$. What we have shown is that $A_n \rightarrow A$ in the strong operator topology.

Let B be the closed unit ball in H . Because A is a compact operator, by the Heine-Borel theorem $A(B)$ is totally bounded: If $\epsilon > 0$, then there is some m and $h_1, \dots, h_m \in H$ such that

$$A(B) \subseteq \bigcup_{j=1}^m B_\epsilon(Ah_j).$$

If $h \in B$, there is some $1 \leq j \leq m$ with $Ah \in B_\epsilon(Ah_j)$, i.e. $\|Ah - Ah_j\| < \epsilon$. If $n \geq 1$, then, as $\|P_n\| \leq 1$,

$$\begin{aligned} \|Ah - A_n h\| &\leq \|Ah - Ah_j\| + \|Ah_j - A_n h_j\| + \|A_n h_j - A_n h\| \\ &= \|Ah - Ah_j\| + \|Ah_j - A_n h_j\| + \|P_n(Ah_j - Ah)\| \\ &\leq 2\|Ah - Ah_j\| + \|Ah_j - A_n h_j\| \\ &\leq 2\epsilon + \|Ah_j - A_n h_j\|. \end{aligned}$$

As $A_n \rightarrow A$ in the strong operator topology, for each $1 \leq j \leq m$ there is some $N(j)$ such that if $n \geq N(j)$ then $\|Ah_j - A_n h_j\| < \epsilon$ and so, if $n \geq N(j)$,

$$\|Ah - A_n h\| < 3\epsilon.$$

Let $N = \max_{1 \leq j \leq m} N(j)$, whence for all $h \in B$, if $n \geq N$ then

$$\|Ah - A_n h\| < 3\epsilon.$$

But

$$\|A - A_n\| = \sup_{\|h\| \leq 1} \|(A - A_n)h\|,$$

so if $n \geq N$ then

$$\|A - A_n\| < 3\epsilon,$$

showing that $A_n \rightarrow A$. □

5 Diagonalizable operators

If \mathcal{E} is an orthonormal set in a Hilbert space H , which we do not demand be separable, then \mathcal{E} is an orthonormal basis for H if and only if for every $h \in H$ we have $h = \sum_{e \in \mathcal{E}} \langle h, e \rangle e$.⁷ In other words, if \mathcal{E} is an orthonormal set in H , then \mathcal{E} is an orthonormal basis for H if and only if

$$\text{id}_H = \sum_{e \in \mathcal{E}} e \otimes e,$$

where the series converges in the strong operator topology.

We say that a linear map $A : H \rightarrow H$ is *diagonalizable* if there is an orthonormal basis of H each element of which is an eigenvector of A . If A is a bounded linear operator on H , A is diagonalizable if and only if there is an orthonormal basis $e_i, i \in I$, of H and $\lambda_i \in \mathbb{C}$ such that the series

$$\sum_{i \in I} \lambda_i e_i \otimes e_i$$

converges to A in the strong operator topology. One checks that the series $\sum_{i \in I} \lambda_i e_i \otimes e_i$ converges to A^* in the strong operator topology.

If $A \in \mathcal{B}(H)$ is diagonalizable with eigenvalues $\{\lambda_i : i \in I\}$, it is a fact that

$$\|A\| = \sup_{i \in I} |\lambda_i|. \tag{1}$$

Theorem 11. *Let H be a Hilbert space with orthonormal basis $\{e_i : i \in I\}$, let $\lambda_i \in \mathbb{C}$, and define a linear map $A : \text{span}\{e_i : i \in I\} \rightarrow \text{span}\{e_i : i \in I\}$ by $Ae_i = \lambda_i e_i$. If $\sup_{i \in I} |\lambda_i| < \infty$, then A extends to a unique element of $\mathcal{B}(H)$.*

Proof. Let $M = \sup_{i \in I} |\lambda_i| < \infty$. If J is a finite subset of I and $x = \sum_{i \in J} \alpha_i e_i$, then, as the e_i are orthonormal,

$$\|Ax\|^2 = \left\| \sum_{i \in J} \alpha_i Ae_i \right\|^2 = \left\| \sum_{i \in J} \alpha_i \lambda_i e_i \right\|^2 = \sum_{i \in J} \|\alpha_i \lambda_i e_i\|^2 \leq M^2 \sum_{i \in J} |\alpha_i|^2.$$

But

$$\|x\|^2 = \left\| \sum_{i \in J} \alpha_i e_i \right\|^2 = \sum_{i \in J} \|\alpha_i e_i\|^2 = \sum_{i \in J} |\alpha_i|^2.$$

So

$$\|Ax\| \leq M\|x\|.$$

It follows that A is a bounded operator on $\text{span}\{e_i : i \in I\}$, which is dense in H . Then there is a unique element of $\mathcal{B}(H)$ whose restriction to $\text{span}\{e_i : i \in I\}$ is equal to A , and we denote this element of $\mathcal{B}(H)$ by A . □

⁷John B. Conway, *A Course in Functional Analysis*, second ed., p. 16, Theorem 4.13.

In Theorem 8 we proved that the image of a compact operator is separable. Hence if a compact operator is diagonalizable then it has only countably many nonzero eigenvalues.

Theorem 12. *If H is a separable Hilbert space with orthonormal basis $e_n, n \geq 1$, and if $A : H \rightarrow H$ is linear and $Ae_n = \lambda_n e_n$ for all n , then $A \in \mathcal{B}_0(H)$ if and only if $\lambda_n \rightarrow 0$ as $n \rightarrow \infty$.*

Proof. Suppose that A is compact. Then A^* is compact, and as A is diagonalizable so is A^* . The proof of Theorem 10 wasn't generally useful enough to be worth putting into a lemma, but to understand the following it will be necessary to read that proof. Let $\lambda_{a(n)}$ be the nonzero eigenvalues of A , and let $\mu_n = \lambda_{a(n)}$, $f_n = e_{a(n)}$. Check that $f_n, n \geq 1$ is an orthonormal basis for $\overline{A^*(H)}$, P_n be the projection onto $\bigvee_{j=1}^n \{f_j\}$. Then using the argument in the proof of Theorem 10 we get $P_n A^* \rightarrow A^*$; this completes our trip back to that proof. Let $A_n = A - AP_n$. As $\|P_n A^* - A^*\| \rightarrow 0$ as $n \rightarrow \infty$,

$$\begin{aligned} \|A_n\| &= \|(A - AP_n)^*\| \\ &= \|A^* - P_n^* A^*\| \\ &= \|A^* - P_n A^*\| \\ &\rightarrow 0. \end{aligned}$$

If $1 \leq j \leq n$ then $A_n f_j = A f_j - A f_j = 0$, and if $j > n$ then $A_n f_j = A f_j = \mu_j f_j$. Hence A_n is a diagonalizable operator, and by (1) we get $\|A_n\| = \sup_{j>n} |\mu_j|$. Together with $\|A_n\| \rightarrow 0$ as $n \rightarrow \infty$, this means that $\limsup_{n \rightarrow \infty} |\mu_n| = 0$. As μ_n are precisely the nonzero λ_n , we obtain from this that $\lim_{n \rightarrow \infty} |\lambda_n| = 0$.

On the other hand, suppose that $\lambda_n \rightarrow 0$ as $n \rightarrow \infty$. Because $\lambda_n \rightarrow 0$, the absolute values of the eigenvalues of A are bounded and hence $A \in \mathcal{B}(H)$. Let P_n be the projection onto $\bigvee_{j=1}^n \{e_j\}$. If $1 \leq j \leq n$ then $AP_n e_j = A e_j = \lambda_j e_j$, and if $j > n$ then $AP_n e_j = 0$. Therefore $AP_n \in \mathcal{B}_{00}(H)$. For $A_n = A - AP_n$, we have $\|A_n\| = \sup_{j>n} |\lambda_j|$, from which it follows that

$$\lim_{n \rightarrow \infty} \|A_n\| = 0.$$

Hence $AP_n \rightarrow A$, and as AP_n are bounded finite rank operators, A is a compact operator. \square

If $A \in \mathcal{B}(H)$ is diagonalizable, say $Ae_i = \lambda_i e_i$, $i \in I$, then $A^* e_i = \overline{\lambda_i} e_i$, $i \in I$, and for $j \in I$,

$$AA^* e_j = A \sum_{i \in I} \overline{\lambda_i} \langle e_j, e_i \rangle e_i = A (\overline{\lambda_j} e_j) = \overline{\lambda_j} \lambda_j e_j = |\lambda_j|^2 e_j.$$

Likewise we get $A^* A e_j = |\lambda_j|^2 e_j$, so $AA^* = A^* A$, that is, a bounded diagonalizable operator is *normal*. The following theorem states an implication in the other direction.⁸ This is an instance of the *spectral theorem*.

Theorem 13. *If H is a Hilbert space over \mathbb{C} and T is a normal compact operator on H , then T is diagonalizable.*

⁸Gert K. Pedersen, *Analysis Now*, revised printing, p. 108, Theorem 3.3.8.

6 Polar decomposition

It is a fact that $A \in \mathcal{B}(H)$ is self-adjoint if and only if $\langle Ax, x \rangle \in \mathbb{R}$ for all $x \in H$. $A \in \mathcal{B}(H)$ is said to be *positive* if A is self-adjoint and $\langle Ax, x \rangle \geq 0$ for all $x \in H$. It is a fact that if $A \in \mathcal{B}(H)$ is positive then there is a unique positive element of $\mathcal{B}(H)$, call it $A^{1/2}$, such that $(A^{1/2})^2 = A$; namely, a bounded positive operator has a unique positive square root in $\mathcal{B}(H)$.⁹ It is straightforward to check that if $A \in \mathcal{B}(H)$ then A^*A is positive, and hence has a square root $(A^*A)^{1/2} \in \mathcal{B}(H)$, which we denote by $|A|$. $|A|$ satisfies $\|Ax\| = \||A|x\|$ for all $x \in H$.

An *isometry* from one Hilbert space to another is a linear map $A : H_1 \rightarrow H_2$ such that if $x \in H_1$ then $\|Ax\| = \|x\|$. If $A : H \rightarrow H$ is linear and the restriction of A to $(\ker A)^\perp$ is an isometry, then we say that A is a *partial isometry*. One checks that a partial isometry is an element of $\mathcal{B}(H)$, if it is not the zero map then it has norm 1, and that its image is closed. We call $(\ker A)^\perp$ the *initial space* of A , and $A(H)$ the *final space* of A . We can prove that if A is a partial isometry then A^* is a partial isometry whose initial space is the final space of A and whose final space is the initial space of A . For example, an orthogonal projection is a partial isometry whose initial space is the image of the orthogonal projection and whose final space is equal to its initial space.

It is a fact that if $A \in \mathcal{B}(H)$ then there is a partial isometry U with $\ker U = \ker A$ satisfying $A = U|A|$, and that if V is a partial isometry with $\ker V = \ker A$ that satisfies $A = V|A|$, then $V = U$.¹⁰ $A = U|A|$ is called the *polar decomposition* of A . The polar decomposition satisfies

$$U^*U|A| = |A|, \quad U^*A = |A|, \quad UU^*A = A, \quad A^* = |A|U^*, \quad |A^*| = U|A|U^*. \quad (2)$$

We will use these formulas repeatedly when we are working with trace class operators, and we have numbered the above equation to refer to it and also to draw the eye to it.

If I is an ideal of $\mathcal{B}(H)$ and $A \in I$ then $|A| = U^*A \in I$. In particular, if $A \in \mathcal{B}_{00}(H)$ then $|A| \in \mathcal{B}_{00}(H)$ and if $A \in \mathcal{B}_0(H)$ then $|A| \in \mathcal{B}_0(H)$.

7 Hilbert-Schmidt operators

Let H be a Hilbert space, and $\{e_i : i \in I\}$ an orthonormal basis of H . We say that $A \in \mathcal{B}(H)$ is a *Hilbert-Schmidt* operator if

$$\sum_{i \in I} \|Ae_i\|^2 < \infty.$$

The following theorem shows that if $A \in \mathcal{B}(H)$ is a Hilbert-Schmidt operator using one basis it will also be one using any other basis, that if A is not a Hilbert-Schmidt operator using one basis it will not be one using any other basis, and that A is Hilbert-Schmidt if and only if its adjoint A^* is Hilbert-Schmidt.

⁹Gert K. Pedersen, *Analysis Now*, revised printing, p. 92, Proposition 3.2.11.

¹⁰Gert K. Pedersen, *Analysis Now*, revised printing, p. 96, Theorem 3.2.17.

Theorem 14. Let H be a Hilbert space. If $\{e_\alpha : \alpha \in I\}$ and $\{f_\alpha : \alpha \in J\}$ are orthonormal bases for H and $A \in \mathcal{B}(H)$, then

$$\sum_{\alpha \in I} \|Ae_\alpha\|^2 = \sum_{\alpha \in J} \|A^*f_\alpha\|^2 = \sum_{\alpha \in J} \|Af_\alpha\|^2.$$

Proof. For each $\beta \in J$, using Parseval's identity we have

$$Af_\beta = \sum_{\alpha \in I} \langle Af_\beta, f_\alpha \rangle f_\alpha, \quad \|Af_\beta\|^2 = \sum_{\alpha \in I} |\langle Af_\beta, f_\alpha \rangle|^2.$$

Using this we get

$$\begin{aligned} \sum_{\beta \in J} \|Af_\beta\|^2 &= \sum_{\beta \in J} \sum_{\alpha \in I} |\langle Af_\beta, f_\alpha \rangle|^2 \\ &= \sum_{\alpha \in I} \sum_{\beta \in J} |\langle Af_\beta, f_\alpha \rangle|^2 \\ &= \sum_{\alpha \in I} \sum_{\beta \in J} |\langle f_\beta, A^*f_\alpha \rangle|^2 \\ &= \sum_{\alpha \in I} \sum_{\beta \in J} |\langle A^*f_\alpha, f_\beta \rangle|^2 \\ &= \sum_{\alpha \in I} \|A^*f_\alpha\|^2. \end{aligned}$$

For each $\beta \in I$, Parseval's identity gives us

$$\|Ae_\beta\|^2 = \sum_{\alpha \in J} |\langle Ae_\beta, f_\alpha \rangle|^2,$$

and using this we obtain

$$\begin{aligned} \sum_{\beta \in I} \|Ae_\beta\|^2 &= \sum_{\beta \in I} \sum_{\alpha \in J} |\langle Ae_\beta, f_\alpha \rangle|^2 \\ &= \sum_{\alpha \in J} \sum_{\beta \in I} |\langle Ae_\beta, f_\alpha \rangle|^2 \\ &= \sum_{\alpha \in J} \sum_{\beta \in I} |\langle e_\beta, A^*f_\alpha \rangle|^2 \\ &= \sum_{\alpha \in J} \sum_{\beta \in I} |\langle A^*f_\alpha, e_\beta \rangle|^2 \\ &= \sum_{\alpha \in J} \|A^*f_\alpha\|^2. \end{aligned}$$

□

Let $\mathcal{B}_2(H)$ be the set of Hilbert-Schmidt operators in $\mathcal{B}(H)$. If $\{e_i : i \in I\}$ is an orthonormal basis for H and $A \in \mathcal{B}(H)$, we define

$$\|A\|_2 = \left(\sum_{i \in I} \|Ae_i\|^2 \right)^{1/2}.$$

To say that $A \in \mathcal{B}_2(H)$ is to say that $A \in \mathcal{B}(H)$ and $\|A\|_2 < \infty$. Using the triangle inequality in $\ell^2(I)$, one checks that $\mathcal{B}_2(H)$ is a vector space and that $\|\cdot\|_2$ is a norm on $\mathcal{B}_2(H)$, which we call the *Hilbert-Schmidt norm*.

Because $\| |A|x \| = \|Ax\|$ for all $x \in H$, A is Hilbert-Schmidt if and only if $|A|$ is Hilbert-Schmidt, and $\|A\|_2 = \| |A| \|_2$. From Theorem 14 we obtain that if $A \in \mathcal{B}_2(H)$ then $A^* \in \mathcal{B}_2(H)$, and $\|A\|_2 = \|A^*\|_2$.

Theorem 15. $\mathcal{B}_2(H)$ is a two sided $*$ -ideal in the C^* -algebra $\mathcal{B}(H)$, and if $A \in \mathcal{B}_2(H)$ and $T \in \mathcal{B}(H)$ then

$$\|AT\|_2 \leq \|A\|_2 \cdot \|T\|, \quad \|TA\|_2 \leq \|T\| \cdot \|A\|_2.$$

Proof. Let \mathcal{E} be an orthonormal basis for H . If $A \in \mathcal{B}_2(H)$ and $T \in \mathcal{B}(H)$, then

$$\|TA\|_2^2 = \sum_{e \in \mathcal{E}} \|TAe\|^2 \leq \sum_{e \in \mathcal{E}} (\|T\| \|Ae\|)^2 = \|T\|^2 \|A\|_2^2 < \infty.$$

On the other hand, using the above and the fact that if A is Hilbert-Schmidt then A^* is Hilbert-Schmidt,

$$\|AT\|_2^2 = \|(AT)^*\|_2^2 = \|T^*A^*\|_2^2 = \|T^*\|^2 \|A^*\|_2^2 < \infty.$$

□

The following theorem shows that the operator norm is dominated by the Hilbert-Schmidt norm, and therefore that the topology on the normed space $\mathcal{B}_2(H)$ with the Hilbert-Schmidt norm is finer than the subspace topology it inherits from $\mathcal{B}(H)$ (i.e. its topology as a normed space with the operator norm).

Theorem 16. If $A \in \mathcal{B}_2(H)$ then $\|A\| \leq \|A\|_2$.

Proof. Let $\epsilon > 0$. We have

$$\|A\| = \sup_{\|v\|=1} \|Av\|.$$

Take $v \in H$, $\|v\| = 1$, with $\|Av\|^2 + \epsilon > \|A\|^2$. There is an orthonormal basis $\{e_\alpha : \alpha \in I\}$ for H that includes v ; one proves this using Zorn's lemma. Then

$$\|A\|^2 < \epsilon + \|Av\|^2 \leq \epsilon + \|Av\|^2 + \sum_{e_\alpha \neq v} \|Ae_\alpha\|^2 = \epsilon + \sum_{\alpha \in I} \|Ae_\alpha\|^2 = \epsilon + \|A\|_2^2.$$

As this is true for all $\epsilon > 0$, it follows that $\|A\| \leq \|A\|_2$. □

The following theorem states that every bounded finite rank operator is a bounded Hilbert-Schmidt operator, and that every bounded Hilbert-Schmidt operator is the limit in the Hilbert-Schmidt norm of a sequence of bounded finite rank operators.

Theorem 17. $\mathcal{B}_{00}(H)$ is a dense subset of the normed space $\mathcal{B}_2(H)$ with the Hilbert-Schmidt norm.

Proof. If $A \in \mathcal{B}_{00}(H)$, then there is an orthonormal basis $\{e_i : i \in I\}$ for H and a finite subset J of I such that if $i \in I \setminus J$ then $Ae_i = 0$. From this it follows that $\|A\|_2 < \infty$, and thus $A \in \mathcal{B}_2(H)$, so $\mathcal{B}_{00}(H)$ is a subset of $\mathcal{B}_2(H)$.

Let $\{e_i : i \in I\}$ be an orthonormal basis for H , let $A \in \mathcal{B}_2(H)$, and let $\epsilon > 0$. As $\sum_{i \in I} \|Ae_i\|^2 < \infty$, there is some finite subset J of I such that

$$\sum_{i \in I \setminus J} \|Ae_i\|^2 < \epsilon.$$

Let P be the orthogonal projection onto $\text{span}\{e_i : i \in J\}$ (which is finite dimensional and hence closed), and define $B \in \mathcal{B}_{00}(H)$ by $B = AP$. We have

$$\|A - B\|_2^2 = \sum_{i \in I \setminus J} \|Ae_i\|^2 < \epsilon,$$

showing that $\mathcal{B}_{00}(H)$ is dense in $\mathcal{B}_2(H)$. □

If $A \in \mathcal{B}_2(H)$, then by the above theorem there is a sequence of bounded finite rank operators A_n such that $\|A_n - A\|_2 \rightarrow 0$ as $n \rightarrow \infty$. But by Theorem 16, $\|A_n - A\| \leq \|A_n - A\|_2$, and by Theorem 6, a limit of bounded finite rank operators is a compact operator, so A is compact. Thus, a bounded Hilbert-Schmidt operator is a compact operator.

We are going to define an inner product on $\mathcal{B}_2(H)$ and we will show that with this inner product $\mathcal{B}_2(H)$ is a Hilbert space. However the cleanest way I know to do this is by defining the trace of an operator. Moreover, we care just as much about trace class operators as we do Hilbert-Schmidt operators.

8 Trace class operators

If $\{e_\alpha : \alpha \in I\}$ and $\{f_\alpha : \alpha \in J\}$ are orthonormal bases for H and $A \in \mathcal{B}(H)$, then using Theorem 14 we have

$$\begin{aligned}
 \sum_{\alpha \in I} \langle |A|e_\alpha, e_\alpha \rangle &= \sum_{\alpha \in I} \langle |A|^{1/2}e_\alpha, |A|^{1/2}e_\alpha \rangle \\
 &= \sum_{\alpha \in I} \| |A|^{1/2}e_\alpha \|^2 \\
 &= \sum_{\alpha \in J} \| |A|^{1/2}f_\alpha \|^2 \\
 &= \sum_{\alpha \in J} \langle |A|^{1/2}f_\alpha, |A|^{1/2}f_\alpha \rangle \\
 &= \sum_{\alpha \in J} \langle |A|f_\alpha, f_\alpha \rangle.
 \end{aligned}$$

If $\{e_i : i \in I\}$ is an orthonormal basis for H , we say that $A \in \mathcal{B}(H)$ is *trace class* if

$$\sum_{i \in I} \langle |A|e_i, e_i \rangle < \infty.$$

We denote the set of trace class operators in $\mathcal{B}(H)$ by $\mathcal{B}_1(H)$. For $A \in \mathcal{B}(H)$, define

$$\|A\|_1 = \sum_{i \in I} \langle |A|e_i, e_i \rangle.$$

To say that $A \in \mathcal{B}_1(H)$ is to say that $A \in \mathcal{B}(H)$ and that $\|A\|_1 < \infty$. As $|A|^{1/2}$ is self-adjoint, it is apparent that $\|A\|_1 = \| |A|^{1/2} \|_2^2$. We will prove that $\mathcal{B}_1(H)$ is a vector space and that $\|\cdot\|_1$ is a norm on this vector space, but this takes a surprising amount of work and we will not do this yet.

The following theorem gives different characterizations of bounded trace class operators.¹¹ This theorem shows in particular that if $A \in \mathcal{B}_1(H)$ then A is the product of two bounded Hilbert-Schmidt operators, and thus, as $\mathcal{B}_2(H)$ is an ideal in $\mathcal{B}(H)$, that $A \in \mathcal{B}_2(H)$. In particular, as a consequence Theorem 16, every bounded Hilbert-Schmidt operator is compact, so every bounded trace class operator is compact.

Theorem 18. *If $A \in \mathcal{B}(H)$, then the following are equivalent.*

- $A \in \mathcal{B}_1(H)$.
- $|A|^{1/2} \in \mathcal{B}_2(H)$.
- A is the product of two elements of $\mathcal{B}_2(H)$.
- $|A|$ is the product of two elements of $\mathcal{B}_2(H)$.

¹¹John B. Conway, *A Course in Operator Theory*, p. 88, Proposition 18.8.

Proof. Let $e_i, i \in I$ be an orthonormal basis for H . Suppose that $A \in \mathcal{B}_1(H)$. We have

$$\| |A|^{1/2} \|_2^2 = \sum_{i \in I} \| |A|^{1/2} e_i \|^2 = \sum_{i \in I} \langle |A|^{1/2} e_i, |A|^{1/2} e_i \rangle = \sum_{i \in I} \langle |A| e_i, e_i \rangle = \|A\|_1,$$

so $|A|^{1/2} \in \mathcal{B}_2(H)$.

Suppose that $|A|^{1/2} \in \mathcal{B}_2(H)$. $A = U|A| = (U|A|^{1/2})|A|^{1/2}$. As $\mathcal{B}_2(H)$ is an ideal, we get $U|A|^{1/2} \in \mathcal{B}_2(H)$, hence A is the product of two elements of $\mathcal{B}_2(H)$.

Suppose that $A = BC$, with $B, C \in \mathcal{B}_2(H)$. Let $A = U|A|$ be the polar decomposition of A . By (2), the polar decomposition satisfies $U^*U|A| = |A|$. But $U|A| = BC$ implies that $U^*U|A| = U^*BC$, hence $|A| = U^*BC = (U^*B)C$. As $\mathcal{B}_2(H)$ is an ideal, we have $U^*B \in \mathcal{B}_2(H)$, so we have written $|A|$ as a product of two elements of $\mathcal{B}_2(H)$.

Suppose that $|A| = BC$, with $B, C \in \mathcal{B}_2(H)$; so $B^* \in \mathcal{B}_2(H)$ too. We have, using the Cauchy-Schwarz inequality first in H and next in $\ell^2(I)$,

$$\begin{aligned} \|A\|_1 &= \sum_{i \in I} \langle |A| e_i, e_i \rangle \\ &= \sum_{i \in I} \langle BC e_i, e_i \rangle \\ &= \sum_{i \in I} \langle C e_i, B^* e_i \rangle \\ &\leq \sum_{i \in I} \|C e_i\| \|B^* e_i\| \\ &\leq \left(\sum_{i \in I} \|C e_i\|^2 \right)^{1/2} \left(\sum_{i \in I} \|B^* e_i\|^2 \right)^{1/2} \\ &= \|C\|_2 \|B^*\|_2 \\ &< \infty. \end{aligned}$$

Hence $A \in \mathcal{B}_1(H)$, completing the proof. \square

The following theorem shows that if $A \in \mathcal{B}_1(H)$ then sums similar to $\|A\|_1$ are also finite, and that the series $\sum_{e \in \mathcal{E}} \langle Ae, e \rangle$ does not depend on the orthonormal basis \mathcal{E} .¹²

Theorem 19. *If $A \in \mathcal{B}_1(H)$ and \mathcal{E} is an orthonormal basis for H , then*

$$\sum_{e \in \mathcal{E}} |\langle Ae, e \rangle| < \infty,$$

and if \mathcal{F} is an orthonormal basis for H then

$$\sum_{e \in \mathcal{E}} \langle Ae, e \rangle = \sum_{f \in \mathcal{F}} \langle Af, f \rangle.$$

¹²John B. Conway, *A Course in Operator Theory*, p. 88, Proposition 18.9.

Proof. By Theorem 18, there are $B, C \in \mathcal{B}_2(H)$ such that $A = C^*B$. If $\lambda \in \mathbb{C}$ and $e \in \mathcal{E}$ then

$$\begin{aligned} \|(B - \lambda C)e\|^2 &= \langle (B - \lambda C)e, (B - \lambda C)e \rangle \\ &= \langle Be, Be \rangle - \langle Be, \lambda Ce \rangle - \langle \lambda Ce, Be \rangle + \langle \lambda Ce, \lambda Ce \rangle \\ &= \|Be\|^2 - \langle Be, \lambda Ce \rangle - \overline{\langle Be, \lambda Ce \rangle} + |\lambda|^2 \|Ce\|^2 \\ &= \|Be\|^2 - 2\operatorname{Re} \langle Be, \lambda Ce \rangle + |\lambda|^2 \|Ce\|^2. \end{aligned}$$

As $\|(B - \lambda C)e\|^2 \geq 0$,

$$2\operatorname{Re} \langle Be, \lambda Ce \rangle \leq \|Be\|^2 + |\lambda|^2 \|Ce\|^2,$$

so

$$2\operatorname{Re} (\overline{\lambda} \langle Be, Ce \rangle) \leq \|Be\|^2 + |\lambda|^2 \|Ce\|^2.$$

This is true for any $\lambda \in \mathbb{C}$ and $e \in \mathcal{E}$. Take $|\lambda| = 1$, depending on e , such that

$$\overline{\lambda} \langle Be, Ce \rangle = |\langle Be, Ce \rangle|,$$

which gives

$$2\operatorname{Re} |\langle Be, Ce \rangle| \leq \|Be\|^2 + \|Ce\|^2,$$

i.e.,

$$|\langle Be, Ce \rangle| \leq \frac{1}{2} (\|Be\|^2 + \|Ce\|^2).$$

Since this inequality doesn't involve λ , the fact that we chose λ depending on e doesn't matter, and the above inequality holds for any $e \in \mathcal{E}$. Therefore

$$\sum_{e \in \mathcal{E}} |\langle Ae, e \rangle| = \sum_{e \in \mathcal{E}} |\langle C^*Be, e \rangle| = \sum_{e \in \mathcal{E}} |\langle Be, Ce \rangle| \leq \frac{1}{2} \|B\|_2^2 + \frac{1}{2} \|C\|_2^2 < \infty,$$

which is the first statement we wanted to prove and which was necessary to prove even to make sense of the second statement.

Because $\sum_{e \in \mathcal{E}} |\langle Ae, e \rangle| < \infty$, the series $\sum_{e \in \mathcal{E}} \langle Ae, e \rangle$ converges. We have to show that its value does not depend on the orthonormal basis \mathcal{E} . If $e \in \mathcal{E}$, then

$$\begin{aligned} \|(B + C)e\|^2 - \|(B - C)e\|^2 &= \|Be\|^2 + \langle Be, Ce \rangle + \langle Ce, Be \rangle + \|Ce\|^2 \\ &\quad - (\|Be\|^2 - \langle Be, Ce \rangle - \langle Ce, Be \rangle + \|Ce\|^2) \\ &= 2\langle Be, Ce \rangle + 2\langle Ce, Be \rangle \\ &= 4\operatorname{Re} \langle Be, Ce \rangle, \end{aligned}$$

which gives us

$$\begin{aligned} \sum_{e \in \mathcal{E}} \operatorname{Re} \langle Ae, e \rangle &= \sum_{e \in \mathcal{E}} \operatorname{Re} \langle Be, Ce \rangle \\ &= \frac{1}{4} \sum_{e \in \mathcal{E}} (\|(B + C)e\|^2 - \|(B - C)e\|^2) \\ &= \frac{1}{4} \|B + C\|_2^2 - \frac{1}{4} \|B - C\|_2^2. \end{aligned}$$

Applying this to $iA = iC^*B$ gives us, as $(iC^*)^* = -iC$,

$$\sum_{e \in \mathcal{E}} \operatorname{Re} \langle iAe, e \rangle = \frac{1}{4} \|B - iC\|_2^2 - \frac{1}{4} \|B + iC\|_2^2.$$

But $\operatorname{Re} \langle iAe, e \rangle = \operatorname{Re} (i \langle Ae, e \rangle) = -\operatorname{Im} \langle Ae, e \rangle$, as $\operatorname{Re} (i(a + ib)) = \operatorname{Re} (ia - b) = -b$. Therefore

$$\sum_{e \in \mathcal{E}} -\operatorname{Im} \langle Ae, e \rangle = \frac{1}{4} \|B - iC\|_2^2 - \frac{1}{4} \|B + iC\|_2^2,$$

i.e.

$$\sum_{e \in \mathcal{E}} \operatorname{Im} \langle Ae, e \rangle = \frac{1}{4} \|B + iC\|_2^2 - \frac{1}{4} \|B - iC\|_2^2.$$

Thus

$$\sum_{e \in \mathcal{E}} \langle Ae, e \rangle = \frac{1}{4} \|B + C\|_2^2 - \frac{1}{4} \|B - C\|_2^2 + \frac{i}{4} \|B + iC\|_2^2 - \frac{i}{4} \|B - iC\|_2^2,$$

and the right-hand side does not depend on the orthonormal basis \mathcal{E} , completing the proof. \square

If $A \in \mathcal{B}_1(H)$ and \mathcal{E} is an orthonormal basis for H , we define the *trace* of A , written $\operatorname{tr} A$, to be

$$\operatorname{tr} A = \sum_{e \in \mathcal{E}} \langle Ae, e \rangle.$$

It is apparent that $\operatorname{tr} : \mathcal{B}_1(H) \rightarrow \mathbb{C}$ is a *positive linear functional*: tr is a linear functional $\mathcal{B}_1(H) \rightarrow \mathbb{C}$, and if $A \in \mathcal{B}_1(H)$ is a positive operator, then $\operatorname{tr} A$ is real and ≥ 0 . If $A \in \mathcal{B}_1(H)$ is a positive operator then it is diagonalizable (being a bounded trace class operator implies that it is compact): there is an orthonormal basis $\{e_i : i \in I\}$ for H such that

$$A = \sum_{i \in I} \langle Ae_i, e_i \rangle e_i \otimes e_i,$$

where the series converges in the strong operator topology. Since A is positive, $\langle Ae_i, e_i \rangle$ is a real nonnegative number for each $i \in I$. $\operatorname{tr} A = 0$ means that

$$\sum_{i \in I} \langle Ae_i, e_i \rangle = 0,$$

and as this is a series of nonnegative terms they must all be 0. Putting these into the expression for A gives us $A = 0$, showing that $\operatorname{tr} : \mathcal{B}_1(H) \rightarrow \mathbb{C}$ is a *positive definite* linear functional. We haven't yet proved that tr is a bounded linear functional. This follows from Theorem 23, which we prove later in this section.

In the following we show that the set of bounded trace class operators is a normed vector space with norm $\|\cdot\|_1$, which we call the *trace norm*.¹³

¹³John B. Conway, *A Course in Operator Theory*, p. 89, Theorem 18.11 (a).

Theorem 20. $\mathcal{B}_1(H)$ is a normed vector space with the norm $\|\cdot\|_1$.

Proof. Let $A, B \in \mathcal{B}_1(H)$, and let their polar decompositions be $A = U|A|$, $B = V|B|$, and $A + B = W|A + B|$. As bounded trace class operators are compact and as the compact operators are a vector space, $A + B$ is a compact operator. We have already stated that if T is compact then $|T|$ is compact, which follows from the polar decomposition of T and the fact that the compact operators are an ideal. Thus $|A + B|$ is a compact operator. $|A + B|$ is a positive operator, so by the spectral theorem for normal compact operators that we stated as Theorem 13, $|A + B|$ is diagonalizable: there is a countable subset $\{e_n : n \geq 1\}$ of an orthonormal basis \mathcal{E} for H , and $\lambda_n \in \mathbb{C}$, such that $|A + B|e_n = \lambda_n e_n$ and $|A + B|e = 0$ if $e \in \mathcal{E}$ is not a member of this countable subset. As $|A + B|$ is a positive operator, the eigenvalues λ_n are real and nonnegative. We have, in the strong operator topology,

$$|A + B| = \sum_{n=1}^{\infty} \lambda_n e_n \otimes e_n.$$

First, as $|A + B| = W^*(A + B)$ and using the Cauchy-Schwarz inequality in H ,

$$\begin{aligned} \|A + B\|_1 &= \sum_{e \in \mathcal{E}} \langle |A + B|e, e \rangle \\ &= \sum_{n \geq 1} \langle |A + B|e_n, e_n \rangle \\ &= \sum_{n \geq 1} \langle (A + B)e_n, W e_n \rangle \\ &= \sum_{n \geq 1} \langle U|A|e_n, W e_n \rangle + \langle V|B|e_n, W e_n \rangle \\ &= \sum_{n \geq 1} \langle |A|e_n, U^* W e_n \rangle + \langle |B|e_n, V^* W e_n \rangle \\ &= \sum_{n \geq 1} \langle |A|^{1/2} e_n, |A|^{1/2} U^* W e_n \rangle + \langle |B|^{1/2} e_n, |B|^{1/2} V^* W e_n \rangle \\ &\leq \sum_{n \geq 1} \| |A|^{1/2} e_n \| \cdot \| |A|^{1/2} U^* W e_n \| + \| |B|^{1/2} e_n \| \cdot \| |B|^{1/2} V^* W e_n \|. \end{aligned}$$

Applying the Cauchy-Schwarz inequality in ℓ^2 to this gives

$$\begin{aligned}
\cdots &\leq \left(\sum_{n \geq 1} \| |A|^{1/2} e_n \|^2 \right)^{1/2} \left(\sum_{n \geq 1} \| |A|^{1/2} U^* W e_n \|^2 \right)^{1/2} \\
&\quad + \left(\sum_{n \geq 1} \| |B|^{1/2} e_n \|^2 \right)^{1/2} \left(\sum_{n \geq 1} \| |B|^{1/2} V^* W e_n \|^2 \right)^{1/2}. \\
&\leq \left(\sum_{e \in \mathcal{E}} \| |A|^{1/2} e \|^2 \right)^{1/2} \left(\sum_{e \in \mathcal{E}} \| |A|^{1/2} U^* W e \|^2 \right)^{1/2} \\
&\quad + \left(\sum_{e \in \mathcal{E}} \| |B|^{1/2} e \|^2 \right)^{1/2} \left(\sum_{e \in \mathcal{E}} \| |B|^{1/2} V^* W e \|^2 \right)^{1/2}.
\end{aligned}$$

So we have

$$\|A + B\|_1 \leq \| |A|^{1/2} \|_2 \cdot \| |A|^{1/2} U^* W \|_2 + \| |B|^{1/2} \|_2 \cdot \| |B|^{1/2} V^* W \|_2.$$

By Theorem 18, $|A|^{1/2}, |B|^{1/2} \in \mathcal{B}_2(H)$, and then using the inequality in Theorem 15 gives us

$$\| |A|^{1/2} U^* W \|_2 \leq \| |A|^{1/2} \|_2 \cdot \|U^*\| \cdot \|W\| \leq \| |A|^{1/2} \|_2,$$

where we used the fact that U^* and W are partial isometries and hence either have norm 1 or 0, depending on whether they are the zero map. Likewise,

$$\| |B|^{1/2} V^* W \|_2 \leq \| |B|^{1/2} \|_2.$$

Therefore we have obtained

$$\|A + B\|_1 \leq \| |A|^{1/2} \|_2^2 + \| |B|^{1/2} \|_2^2 = \|A\|_1 + \|B\|_1.$$

so $A + B \in \mathcal{B}_1(H)$, and $\|\cdot\|_1$ satisfies the triangle inequality.

If $A \in \mathcal{B}(H)$ and $\alpha \in \mathbb{C}$ then $|\alpha A|^{1/2} = |\alpha|^{1/2} |A|^{1/2}$, and from this it follows that if $A \in \mathcal{B}_1(H)$ and $\alpha \in \mathbb{C}$ then $\|\alpha A\|_1 = |\alpha| \|A\|_1$ and so $\alpha A \in \mathcal{B}_1(H)$. Therefore $\mathcal{B}_1(H)$ is a vector space.

If $\|A\|_1 = 0$, then $\text{tr } |A| = 0$. We have shown that tr is a positive definite linear functional: hence $|A| = 0$, and so $A = 0$. This completes the proof that $\|\cdot\|_1$ is a norm on $\mathcal{B}_1(H)$. \square

We have shown that $\mathcal{B}_1(H)$ with the trace norm $\|\cdot\|_1$ is a normed space. In the following theorem we show that a bounded finite rank operator is a bounded trace class operator, and that $\mathcal{B}_{00}(H)$ is a dense subset of $\mathcal{B}_1(H)$.

Theorem 21. $\mathcal{B}_{00}(H)$ is a dense subset of the normed space $\mathcal{B}_1(H)$ with the trace norm.

Proof. If $A \in \mathcal{B}_{00}(H)$, then there is an orthonormal basis $\{e_i : i \in I\}$ for H and a finite subset J of I such that if $i \in I \setminus J$ then $Ae_i = 0$, and so $|A|e_i = 0$. This gives $\|A\|_1 < \infty$, so $A \in \mathcal{B}_1(H)$.

Let $\{e_i : i \in I\}$ be an orthonormal basis for H . We've shown that the bounded finite rank operators are contained in the bounded trace class operators, and now we have to show that if $A \in \mathcal{B}_1(H)$ and $\epsilon > 0$ then there is some $B \in \mathcal{B}_{00}(H)$ satisfying $\|A - B\|_1 < \epsilon$. As $\sum_{i \in I} \langle |A|e_i, e_i \rangle < \infty$, there is a finite subset J of I such that

$$\sum_{i \in I \setminus J} \langle |A|e_i, e_i \rangle < \epsilon.$$

Let P be the orthogonal projection onto $\text{span}\{e_i : i \in J\}$, and define $B \in \mathcal{B}_{00}(H)$ by $B = AP$, which gives us

$$\|A - B\|_1 = \sum_{i \in I} \langle |A - B|e_i, e_i \rangle = \sum_{i \in I \setminus J} \langle |A|e_i, e_i \rangle < \epsilon.$$

□

Taking an adjoint of an operator on a Hilbert space and taking the complex conjugate of a complex number ought to be interchangeable where it makes sense. We show in the following theorem that the adjoint of an element of $\mathcal{B}_1(H)$ is also an element of $\mathcal{B}_1(H)$ and that the trace of the adjoint is the complex conjugate of the trace.

Theorem 22. *If $A \in \mathcal{B}_1(H)$ then $A^* \in \mathcal{B}_1(H)$ and*

$$\text{tr } A^* = \overline{\text{tr } A}.$$

Proof. As $A \in \mathcal{B}_1(H)$, by Theorem 18 there are $B, C \in \mathcal{B}_2(H)$ such that $A = C^*B$. Then $A^* = B^*C$ is a product of two bounded Hilbert-Schmidt operators, and so by the same theorem is itself an element of $\mathcal{B}_1(H)$.

We're going to extract something from our proof of Theorem 19, which showed that the trace of an operator does not depend on the orthonormal basis that we use: We proved that, with $A = C^*B$,

$$\text{tr}(A) = \frac{1}{4}\|B + C\|_2^2 - \frac{1}{4}\|B - C\|_2^2 + \frac{i}{4}\|B + iC\|_2^2 - \frac{i}{4}\|B - iC\|_2^2,$$

Applying this to the adjoint $A^* = B^*C$, and as $i(C + iB) = iC - B$,

$$\begin{aligned} \text{tr } A^* &= \frac{1}{4}\|C + B\|_2^2 - \frac{1}{4}\|C - B\|_2^2 + \frac{i}{4}\|C + iB\|_2^2 - \frac{i}{4}\|C - iB\|_2^2 \\ &= \frac{1}{4}\|B + C\|_2^2 - \frac{1}{4}\|B - C\|_2^2 + \frac{i}{4}\|iC - B\|_2^2 - \frac{i}{4}\|iB + C\|_2^2 \\ &= \overline{\text{tr } A}. \end{aligned}$$

□

It is familiar to us that if A and B are matrices then $\operatorname{tr}(AB) = \operatorname{tr}(BA)$. In the next theorem we show that this is true for bounded linear operators providing one of the two is a bounded trace class operator.¹⁴ The first thing we prove in this theorem is that $\mathcal{B}_1(H)$ is an ideal of the algebra $\mathcal{B}(H)$, so that it makes sense to talk about the trace of a product of two bounded linear operators only one of which is a bounded trace class operator. The theorem also shows that $\operatorname{tr} : \mathcal{B}_1(H) \rightarrow \mathbb{C}$, which we have already shown is a positive definite linear functional, is bounded.

Theorem 23. *If $A \in \mathcal{B}_1(H)$ and $T \in \mathcal{B}(H)$, then $AT, TA \in \mathcal{B}_1(H)$, and $\operatorname{tr}(AT) = \operatorname{tr}(TA)$, and $|\operatorname{tr}(TA)| \leq \|T\| \|A\|_1$.*

Proof. A is the product of two bounded Hilbert-Schmidt operators, say $A = C^*B$ (we write it this way because this will be handy later in the proof). Hence $AT = C^*(BT)$. As $\mathcal{B}_2(H)$ is an ideal of $\mathcal{B}(H)$, we have $BT \in \mathcal{B}_2(H)$, showing that AT is a product of two bounded Hilbert-Schmidt operators, which implies that $AT \in \mathcal{B}_1(H)$. Similarly, $TA \in \mathcal{B}_1(H)$. We use from the proof of Theorem 19 the following: (we wrote $A = C^*B$ to match the way we wrote A in that theorem)

$$\operatorname{tr}(C^*B) = \frac{1}{4}\|B+C\|_2^2 - \frac{1}{4}\|B-C\|_2^2 + \frac{i}{4}\|B+iC\|_2^2 - \frac{i}{4}\|B-iC\|_2^2.$$

Applying this to CB^* , and using that the norm of the adjoint of an operator is equal to the norm of the operator itself and that $(iC^*)^* = -iC$,

$$\begin{aligned} \operatorname{tr}(CB^*) &= \frac{1}{4}\|B^*+C^*\|_2^2 - \frac{1}{4}\|B^*-C^*\|_2^2 + \frac{i}{4}\|B^*+iC^*\|_2^2 - \frac{i}{4}\|B^*-iC^*\|_2^2 \\ &= \frac{1}{4}\|B+C\|_2^2 - \frac{1}{4}\|B-C\|_2^2 + \frac{i}{4}\|B-iC\|_2^2 - \frac{i}{4}\|B+iC\|_2^2 \\ &= \overline{\operatorname{tr}(C^*B)}. \end{aligned}$$

Using this we obtain

$$\begin{aligned} \operatorname{tr}(TA) &= \operatorname{tr}((TC^*)B) \\ &= \overline{\operatorname{tr}((TC^*)^*B^*)} \\ &= \overline{\operatorname{tr}(CT^*B^*)} \\ &= \overline{\operatorname{tr}(C(BT)^*)} \\ &= \operatorname{tr}(C^*(BT)) \\ &= \operatorname{tr}(AT), \end{aligned}$$

which we wanted to show.

We still have to prove that $|\operatorname{tr}(TA)| \leq \|T\| \|A\|_1$. Let $A = U|A|$ be the polar decomposition of A , and let \mathcal{E} be an orthonormal basis for H . By Theorem 18,

¹⁴John B. Conway, *A Course in Operator Theory*, p. 89, Theorem 18.11 (e).

$|A|^{1/2} \in \mathcal{B}_2(H)$. (We mention this to justify talking about the Hilbert-Schmidt norm of $|A|^{1/2}$.) We have, using the Cauchy-Schwarz inequality in H and in ℓ^2 ,

$$\begin{aligned}
|\operatorname{tr}(TA)| &= \left| \sum_{e \in \mathcal{E}} \langle TU|A|e, e \rangle \right| \\
&= \left| \sum_{e \in \mathcal{E}} \langle |A|^{1/2}e, |A|^{1/2}U^*T^*e \rangle \right| \\
&\leq \sum_{e \in \mathcal{E}} |\langle |A|^{1/2}e, |A|^{1/2}U^*T^*e \rangle| \\
&\leq \sum_{e \in \mathcal{E}} \| |A|^{1/2}e \| \cdot \| |A|^{1/2}U^*T^*e \| \\
&\leq \left(\sum_{e \in \mathcal{E}} \| |A|^{1/2}e \|^2 \right)^{1/2} \left(\sum_{e \in \mathcal{E}} \| |A|^{1/2}U^*T^*e \|^2 \right)^{1/2} \\
&= \| |A|^{1/2} \|_2 \cdot \| |A|^{1/2}U^*T^* \|_2.
\end{aligned}$$

By Theorem 15, and as U^* is a partial isometry,

$$\| |A|^{1/2}U^*T^* \|_2 \leq \| |A|^{1/2} \|_2 \cdot \|U^*\| \cdot \|T^*\| \leq \| |A|^{1/2} \|_2 \cdot \|T^*\|.$$

Therefore

$$|\operatorname{tr}(TA)| \leq \| |A|^{1/2} \|_2^2 \cdot \|T^*\| = \| |A|^{1/2} \|_2^2 \cdot \|T\| = \|A\|_1 \cdot \|T\|,$$

completing the proof. \square

In Theorem 22 we proved that the adjoint of a bounded trace class operator is itself trace class, and we now prove that they have the same trace norm.

Theorem 24. *If $A \in \mathcal{B}_1(H)$ then*

$$\|A^*\|_1 = \|A\|_1.$$

Proof. The polar decomposition $A = U|A|$ satisfies $|A^*| = U|A|U^*$ and $U^*U|A| = |A|$, as we stated in (2). We have, using this and Theorem 23,

$$\|A^*\|_1 = \operatorname{tr} |A^*| = \operatorname{tr} ((U|A|)U^*) = \operatorname{tr} (U^*(U|A|)) = \operatorname{tr} |A| = \|A\|_1.$$

\square

Theorem 25. *If $A \in \mathcal{B}_1(H)$ and $T \in \mathcal{B}(H)$, then*

$$\|AT\|_1 \leq \|A\|_1 \cdot \|T\|, \quad \|TA\|_1 \leq \|T\| \cdot \|A\|_1.$$

Proof. Let A have the polar decomposition $A = U|A|$ and let TA have the polar decomposition $TA = W|TA|$. We have, from (2),

$$|TA| = W^*(TA) = W^*TU|A|.$$

It follows from Theorem 23 that, putting $S = W^*TU$,

$$\|TA\|_1 = \operatorname{tr}(|TA|) = \operatorname{tr}(S|A|) \leq \|S\| \cdot \|A\|_1.$$

As W^* and U are partial isometries, $\|S\| \leq \|T\|$, thus

$$\|TA\|_1 \leq \|T\| \cdot \|A\|_1. \quad (3)$$

On the other hand, Theorem 24 tells us

$$\|AT\|_1 = \|T^*A^*\|_1.$$

As $A^* \in \mathcal{B}_1(H)$, by (3) we get

$$\|T^*A^*\|_1 \leq \|T^*\| \cdot \|A^*\|_1 = \|T\| \cdot \|A\|_1,$$

using Theorem 24 again. \square

In Theorem 16 we proved that if $A \in \mathcal{B}_2(H)$ then $\|A\| \leq \|A\|_2$. Now we prove that the trace norm also dominates the operator norm, so that the topology on the normed space $\mathcal{B}_1(H)$ with the trace norm is finer than its topology as a subspace of $\mathcal{B}(H)$.

Theorem 26. *If $A \in \mathcal{B}_1(H)$, then*

$$\|A\| \leq \|A\|_1.$$

Proof. Let A have polar decomposition $A = U|A|$. As $|A|$ is a compact positive, it is diagonalizable: there is an orthonormal basis $\{e_i : i \in I\}$ for H and $\lambda_i \in \mathbb{C}$ such that $|A|e_i = \lambda_i e_i$. On the one hand, the operator norm of a diagonalizable operator is the supremum of the absolute values of its eigenvalues, as we stated in (1). On the other hand,

$$\| |A| \|_1 = \sum_{i \in I} |\lambda_i|.$$

Certainly then $\| |A| \| \leq \| |A| \|_1$. But

$$\|A\| \leq \|U\| \cdot \| |A| \| \leq \| |A| \| \leq \| |A| \|_1 = \|A\|_1,$$

which is what we wanted to prove. \square

We have already shown that the trace class operators with the trace norm are a normed space. We now prove that they are a Banach space. We do this by showing that there is an isometric isomorphism $\rho : \mathcal{B}_1(H) \rightarrow \mathcal{B}_0(H)^*$.¹⁵ The latter space is a Banach space, so if $A_n \in \mathcal{B}_1(H)$ is a Cauchy sequence, its image $\rho(A_n)$ is a Cauchy sequence in $\mathcal{B}_0(H)^*$ and hence has a limit, call it B . Since ρ is surjective, there is some $A \in \mathcal{B}_1(H)$ such that $\rho(A) = B$, and one checks that $A_n \rightarrow A$.

For $A \in \mathcal{B}_1(H)$ and $C \in \mathcal{B}_0(H)$, define

$$\Phi_A(C) = \text{tr}(CA).$$

Theorem 27. *The map $\rho : \mathcal{B}_1(H) \rightarrow \mathcal{B}_0(H)^*$ defined by*

$$\rho(A) = \Phi_A, \quad A \in \mathcal{B}_1(H),$$

is an isometric isomorphism.

Proof. Let $A \in \mathcal{B}_1(H)$. It is apparent that $\Phi_A : \mathcal{B}_0(H) \rightarrow \mathbb{C}$ is a linear map. Using Theorem 23, its operator norm is

$$\|\Phi_A\| = \sup_{\|C\| \leq 1} |\Phi_A(C)| = \sup_{\|C\| \leq 1} |\text{tr}(CA)| \leq \sup_{\|C\| \leq 1} \|C\| \|A\|_1 = \|A\|_1,$$

where the supremum is taken over compact operators. Hence $\Phi_A \in \mathcal{B}_0(H)^*$ (if $H = \{0\}$ then the final equality is \leq). We have $\|\Phi_A\| \leq \|A\|_1$, so to show that ρ is an isometric isomorphism, we have to show that if $A \in \mathcal{B}_1(H)$ then $\|\Phi_A\| \geq \|A\|_1$, and that ρ is surjective (as being injective is implied by being an isometry).

Let $\Phi \in \mathcal{B}_0(H)^*$. For $g, h \in H$, define

$$B(g, h) = \Phi(g \otimes h), \quad B(g, h)v = g \otimes h(v) = \langle v, h \rangle g.$$

It is apparent from this that B is a sesquilinear form on H . A sesquilinear form B is said to be *bounded* if $M = \sup_{\|g\|, \|h\|=1} |B(g, h)| < \infty$. For $\|g\| = \|h\| = 1$,

$$|B(g, h)| = |\Phi(g \otimes h)| \leq \|\Phi\| \|g \otimes h\| \leq \|\Phi\| \|g\| \|h\| = \|\Phi\|.$$

Thus B is a bounded sesquilinear form on H , and we can therefore apply the *Riesz representation theorem*,¹⁶ which states that there is a unique $T \in \mathcal{B}(H)$ such that

$$B(g, h) = \langle g, Th \rangle, \quad g, h \in H,$$

and that this T satisfies $\|T\| = M$.

Let $A = T^*$, let $A = U|A|$ be the polar decomposition of A , and let \mathcal{E} be an orthonormal basis for H . If \mathcal{E}_0 is a finite subset of \mathcal{E} , define

$$C_{\mathcal{E}_0} = \left(\sum_{e \in \mathcal{E}_0} e \otimes e \right) U^* = \sum_{e \in \mathcal{E}_0} e \otimes Ue.$$

¹⁵John B. Conway, *A Course in Operator Theory*, p. 93, Theorem 19.1.

¹⁶Walter Rudin, *Functional Analysis*, second ed., p. 310, Theorem 12.8.

It is apparent that $C_{\mathcal{E}_0} \in \mathcal{B}_{00}(H)$, and one checks that $\|C_{\mathcal{E}_0}\| \leq 1$. We have

$$\begin{aligned}
\sum_{e \in \mathcal{E}_0} \langle |A|e, e \rangle &= \left| \sum_{e \in \mathcal{E}_0} \langle |A|e, e \rangle \right| \\
&= \left| \sum_{e \in \mathcal{E}_0} \langle U^* A e, e \rangle \right| \\
&= \left| \sum_{e \in \mathcal{E}_0} \langle e, T U e \rangle \right| \\
&= \left| \sum_{e \in \mathcal{E}_0} B(e, U e) \right| \\
&= \left| \sum_{e \in \mathcal{E}_0} \Phi(e \otimes U e) \right| \\
&= |\Phi(C_{\mathcal{E}_0})|.
\end{aligned}$$

Then

$$\sum_{e \in \mathcal{E}_0} \langle |A|e, e \rangle \leq \|\Phi\| \|C_{\mathcal{E}_0}\| \leq \|\Phi\|.$$

This is true for any finite subset \mathcal{E}_0 of \mathcal{E} , and it follows that

$$\|A\|_1 \leq \|\Phi\|, \tag{4}$$

and thus $A \in \mathcal{B}_1(H)$.

Let $C \in \mathcal{B}_{00}(H)$. Then there are some $g_1, \dots, g_n, h_1, \dots, h_n \in H$ such that (cf. Theorem 1)

$$C = \sum_{k=1}^n g_k \otimes h_k.$$

We have

$$\begin{aligned}
\Phi(C) &= \sum_{k=1}^n \Phi(g_k \otimes h_k) \\
&= \sum_{k=1}^n B(g_k, h_k) \\
&= \sum_{k=1}^n \langle A g_k, h_k \rangle \\
&= \sum_{k=1}^n \text{tr}(A(g_k \otimes h_k)) \\
&= \text{tr}(AC) \\
&= \Phi_A(C).
\end{aligned}$$

Since the bounded linear functionals Φ and Φ_A agree on $\mathcal{B}_{00}(H)$, a dense subset of $\mathcal{B}_1(H)$ with the trace norm, they are equal. Therefore, ρ is surjective. Moreover, we showed in (4) that $\|A\|_1 \leq \|\Phi\|$, so $\|\Phi_A\| \geq \|A\|_1$, so ρ is an isometry, which completes the proof. \square

We have shown that the $\mathcal{B}_1(H)$ is the dual of $\mathcal{B}_0(H)$. It can further be shown that $\mathcal{B}(H)$ is the dual of $\mathcal{B}_1(H)$. For $T \in \mathcal{B}(H)$ we define $\Psi_T : \mathcal{B}_1(H) \rightarrow \mathbb{C}$ by $\Psi_T(A) = \text{tr}(TA)$, $A \in \mathcal{B}_1(H)$. Then the map $T \mapsto \Psi_T$ is an isometric isomorphism $\mathcal{B}(H) \rightarrow \mathcal{B}_1(H)^*$.¹⁷

9 The Hilbert-Schmidt inner product

If $A, B \in \mathcal{B}_2(H)$, we define

$$\langle A, B \rangle = \text{tr}(B^*A),$$

which makes sense because, by Theorem 18, $AB \in \mathcal{B}_1(H)$. As $\text{tr} : \mathcal{B}_1(H) \rightarrow \mathbb{C}$ is a positive definite linear functional, we get that $\langle \cdot, \cdot \rangle : \mathcal{B}_2(H) \times \mathcal{B}_2(H) \rightarrow \mathbb{C}$ is an inner product. We call this the *Hilbert-Schmidt inner product*. Check that $\langle A, A \rangle = \|A\|_2^2$. It is a fact that $\mathcal{B}_2(H)$ is a complete metric space with the Hilbert-Schmidt norm,¹⁸ and hence $\mathcal{B}_2(H)$ with the Hilbert-Schmidt inner product is a Hilbert space.

Theorem 28. $\mathcal{B}_2(H)$ with the Hilbert-Schmidt inner product is a Hilbert space.

¹⁷John B. Conway, *A Course in Operator Theory*, p. 94, Theorem 19.2.

¹⁸Gert K. Pedersen, *Analysis Now*, revised printing, p. 119, Theorem 3.4.9.