

Microeconomic Theory I

Midterm

Marcin Peški

October 27, 2015

Each question has the same value. You need to provide arguments for each answer. If you cannot solve one part of the problem, don't give up and try to solve the next one. If the question explicitly asks you to prove a result from the class, you must carefully describe the proof. Otherwise, you may use any result from the class given that you clearly state the assumptions, thesis and verify that the assumptions hold in your application. You have 110 minutes. Good luck!

1. Answer the following questions:

- (a) Suppose that $u : R_+^L \rightarrow R$ is a continuous, locally non-satiated utility function and strictly quasi-concave on consumption set $X = R_+^L$. Derive the Slutsky equation.

- (b) Use the Slutsky equation and Shepherd's Lemma to establish the Law of the Compensated Demand.

(c) Explain that the Law of (Uncompensated) Demand holds for homothetic preferences.

- (d) Let L be the number of inputs, let $w \in R_+^L$ be the vector of input prices, and let $f : R^L \rightarrow R$ be the production function of a firm. Establish the Law of Supply for the firm (using the Hotelling Lemma).

2. In the class, we listed necessary conditions for the Walrasian demand function $x(p, w)$ to be derived as a (possibly, set-) solution to the consumer's problem with monotonic, strictly convex, and continuous preferences. In these question, you will examine whether (some of the) conditions are empirically falsifiable, i.e., whether one can observe an empirical violation of them.

From now on, a *dataset* is a list of finitely many data points (p_i, w_i, x_i) , where $x_i \in R_+^L, p_i \in R_{++}^L, w_i \in R_+$ for each i . Say that a dataset invalidates property P if there is no function $x : R_{++}^L \times R_+ \rightarrow R_+^L$ that satisfies property P and such that for each i , $x(p_i, w_i) = x_i$.

In each of the questions below, start your answer by stating the definition of the relevant property.

- (a) Give an example of dataset that invalidates homogeneity of degree 0 in (p, w) or explain that such a dataset does not exist.

- (b) Give an example of dataset that invalidates Walras Law or explain that such a dataset does not exist.

- (c) Give an example of dataset that invalidates Weak Axiom of Revealed Preferences, but does not violate Walras Law, or explain that such a dataset does not exist. (Hint: It might be helpful to draw a picture.)

- (d) (Harder.) Assume that it is known that $x(p, w)$ satisfies Walras Law, homogeneity and WARP, and it is continuously differentiable in p with a known bound on the derivative. Give an example of a dataset that invalidates the symmetry of the Slutsky matrix, or explain that such a dataset does not exist. (Hint: Why was the symmetry of the Slutsky matrix important for our theory?) To get a credit, describe an idea behind your answer as clearly as you can.

3. Answer the following questions.

(a) State the definition of a supermodular function.

- (b) Suppose that (X, \leq_X) and (Y, \leq_Y) are lattices. Describe a partial order on $X \times Y$ that makes it into a lattice. Carefully define the lattice operations on the product lattice.

- (c) Suppose that (X, \leq_X) is an arbitrary lattice and $Y = R_+$ is a lattice of positive real numbers. Suppose that $f : X \rightarrow R$ is supermodular and increasing. The latter means that for any $x \leq_X x'$, we have $f(x) \leq f(x')$. Show that function $g : X \times Y \rightarrow R$ defined so that

$$g(x, y) = yf(x)$$

is supermodular on the product lattice.

4. Consider a decision maker with Bernoulli utility function $u(\cdot)$ and expected utility preferences over lotteries F .

(a) Define Arrow Pratt risk aversion $r_u(x)$ and certainty equivalent $c(u, F)$ of lottery F .

- (b) Show that for any two increasing utility functions u and v , if $r_u(x) \leq r_v(x)$ for each x , then, for each F

$$c(u, F) \geq c(v, F).$$

Carefully describe the proof of the above claim.

- (c) Recall that a lottery F is a c.d.f. of a probability distribution over prizes z . For each lottery F and each $a \in \mathbb{R}$, let $F + a$ be a lottery obtained from F by adding a to each prize. In other words, $(F + a)(z + a) = F(z)$ for each z . Show that if $r_u(x)$ is strictly decreasing in x , then for each $a > 0$,

$$c(u, F) + a \leq c(u, F + a).$$

- (d) Use the above observation to comment about the change in risk-attitudes for different levels of wealth when the decision maker has (i) Cobb-Douglas utility function $u(c) = c^\alpha$ for some $\alpha \in (0, 1)$, or (ii) the exponential utility $v(x) = -e^{-ax}$.